

- distribution of Amazonian species. *P. Natl. Acad. Sci. (USA)* 97(25):13672–13677.
- Gonzalez-Socoloske, D. and Snarr, K. A. 2010. An incidente de swimming in a large river by a manthed howling monkey (*Alouatta palliata*) on the north coast of Honduras. *Neotrop. Primates*. 17(1):28–31.
- Hershkovitz, P. 1977. *Living New World monkeys (Platyrrhini) with an introduction to primates*. Vol. 1. University of Chicago Press, Chicago.
- Iwanaga, S. and Ferrari, S. F. 2002. Geographic distribution and abundance of woolly (*Lagothrix cana*) and spider (*Ateles chamek*) monkeys in southwestern Brazilian Amazonia. *Am. J. Primatol.* 56:57–64.
- Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V. N., Underwood, E. C., D'Amico, J. A., Itoua, I., Strand, H. E., Morrison, J. C., Loucks, C. J., Allnutt, T. F., Ricketts, T. H., Kura, Y., Lamoreux, J. F., Wettengel, W. W., Hedao, P. and Kassem, K. R. 2001. Terrestrial ecoregions of the world. A new map of life on Earth. *Bioscience*. 51:933–938.
- Parnell, J. P. and Buchanan-Smith, H. M. 2001. An unusual social display by gorillas. *Nature*. 142:294.
- Sick, H. 1967. Rios e enchentes na Amazônia como obstáculo para a avifauna. Manaus, *Atas Simpósio sobre a biota Amazônica*, 5:495–520.
- Souza-Filho, P. W. M., Quadros, M. L. E. S., Scandolaro, J. E., Silva-Filho, E. P. and Reis, M. R. 1999. Compartimentação morfoestrutural e neotectônica do sistema fluvial Guaporé-Mamoré Alto Madeira, Rondônia, Brasil. *Revis. Brasil. Geociên.* 29(4):469–476.
- Wallace, R. B., Painter, R. L. E. and Taber, A. B. 1996. Primate diversity, habitat preferences, and population density estimates in Noel Kempff Mercado National Park, Santa Cruz Department, Bolivia. *Am. J. Phys. Anthropol.* 46(3):197–211.
- Wata, k. 1981. Habitat utilization by wintering Japanese macaques (*Macaca fuscata fuscata*) in the Shiga Heights. *Primates*. 22(3):330–348.
- Yeager, C. P. 1991. Possible antipredator behavior associated with river crossings by proboscis monkeys (*Nasalis larvatus*). *Am. J. Primatol.* 24(1):61–66.

YELLOW-TAILED WOOLLY MONKEY (*OREONAX FLAVICAUDA*: HUMBOLDT 1812) ALTITUDINAL RANGE EXTENSION, UCHIZA, PERÚ

Néstor Allgas
Sam Shanee
Ana Peralta
Noga Shanee

The yellow-tailed woolly monkey (*Oreonax flavicauda*) is endemic to a small area of montane cloud forest dominated by *Ficus* spp. in the Peruvian departments of Amazonas and San Martín (Leo Luna 1980) and neighboring areas of the departments of Huánuco and La Libertad (Graves &

O'Neil, 1980; Parker & Barkley, 1981; Shanee, 2011). This species is listed as Critically Endangered on the IUCN Red List of Threatened Species (2008, A4c) and Endangered on Appendix 1 of CITES (2005). The main threats to *O. flavicauda* are massive deforestation for agriculture, subsistence hunting, logging and mining (deLuycker, 2007; Leo Luna, 1980; Shanee, 2011). In many areas habitat loss has forced this species into small forest fragments (Shanee et al., 2007; Shanee, 2011).

On the 25th and 26th of January 2013, while carrying out distribution surveys of the Andean night monkey (*Aotus miconax*), we encountered a group of *O. flavicauda* 14.5 km west of the city of Uchiza in San Martín department in an area known locally as Tingo de Uchiza (S 8°28'47.04", W 76°35'24.90"), just north of the border with Huánuco (Fig 1.). The group was found along an existing 1.1 km trail at altitudes between 1,084 and 1,373 m. a.s.l., just under 500 m lower than previous observations (Table 1). We observed the group feeding on fruiting figs (*Ficus* spp.) for 25 minutes before they crossed a small stream which feeds the Rio Trisneja where we were unable to follow. The group consisted of 12 individuals, including two females with infants.

The habitat was similar to that described by previous researchers (Shanee, 2011; Shanee & Shanee, 2011) with high humidity (up to 99% relative humidity at 14.7°C). Forests in this area are dominated by Moraceae (*Ficus* spp.) and Cecropiaceae (*Cecropia* spp.) as well as Fabaceae (*Inga* spp. and *Erythrina* spp.), Icacinaceae (*Citronella* spp. and *Styloceras* spp.) with a high density of epiphytes. Our observations were made in a long thin canyon with steep sides that culminated in the 400 m high *Velo de Plata* waterfall. It is possible that the extremely humid and cool microclimate created by the local topography and the effect of the waterfall have allowed the higher altitude forest type, which




Figure 1. Map of observation locality.

Table 1. Minimum altitudes where *O. flavicauda* has been observed in previous studies.

Altitude (m a.s.l.)	Study sites	Source
1,084	Tingo de Uchiza, San Martín	This study
1,560	Shunte, San Martín	Shanee (2011)
> 1,600	Pucatambo, Amazonas	Leo Luna (1980)
1,505	Bosque de Protección Alto Mayo	DeLuyker (2007)
2,400	Ongón, La Libertad	Parker & Barkley (1981)
1,670	Abra Patricia, Amazonas	Graves and O'Neil (1980)
1,550	Pucatambo, Amazonas	Thomas (1927)
2,220	Cordillera de Colán, Amazonas	Butchart et al (1995)

is home to *O. flavicauda*, to establish itself at these lower altitudes.

Local villagers stated that *O. flavicauda* is common in the area, which they had mistakenly identified as howler monkeys (*Alouatta* sp.). When informed about the species endemism and Critically Endangered status they showed a lot of interest in conserving the species. The Municipality of Uchiza is currently working with the San Martin Regional Government to create a new Regional Conservation Area (*Area de Conservación Regional*) which covers the area where our observations were made. More time will need to be spent in this area to see how much lower the species distribution reaches in this and other similar areas. With the current high rates of habitat loss throughout this species distribution area, any additional areas of habitat suitable for them are of importance for its conservation.

References

- Butchart, S. H. M., Barnes, R., Davies, C.W.N., Fernández, M. and Seddon, N., 1995. Observations of two threatened primates in the Peruvian Andes. *Primate Cons.* 16: 15–19.
- DeLuycker, A. M., 2007. Notes on the Yellow-Tailed Woolly Monkey (*Oreonax flavicauda*) and Its Status in the Protected Forest of Alto Mayo, Northern Peru. *Primate Cons.* 22: 41–47.
- Graves, G. R. and O'Neill, J. P., 1980. Notes on the Yellow-Tailed Woolly Monkey (*Lagothrix flavicauda*) of Peru. *J. of Mammal.* 61: 345–357.
- Luna, M. L., 1980. First Field Study of the Yellow-tailed Woolly Monkey. *Oryx* 15: 386–389.
- Parker, T.A. and Barkley, L.J., 1981. New Locality for the Yellow-tailed Woolly Monkey. *Oryx* 16: 71–72.
- Shanee, N., Shanee, S. and Maldonado, A. M., 2007. Conservation assessment and planning for the yellow tailed woolly monkey (*Oreonax flavicauda*) in Peru. *Wildlife Biol. Prac.* 3: 73–82.
- Shanee, S., 2011. Distribution Survey and Threat Assessment of the Yellow-tailed Woolly Monkey (*Oreonax flavicauda*; Humboldt 1812), Northeastern Peru. *Int. J. Primatol.* 32: 691–707.

Shanee, S. and Shanee, N., 2011. Observations of terrestrial behavior in the Peruvian night Monkey (*Aotus miconax*) in an anthropogenic landscape, La Esperanza, Perú. *Neotrop. Primates.* 18: 55–58.

Thomas, O., 1927. A remarkable new monkey from Peru. *Ann. Mag. Nat. Hist.* Ser. 9, 19: 156–157.

Néstor Allgas, Asociación Neotropical Primate Conservation (ANPC) Perú, 1187 Carretera Fernando Belaunde Terry, La Esperanza, Yamborasbamba, Perú y Universidad Nacional Mayor de San Marcos, Facultad de Ciencias Biológicas, Av. Universitaria/Av. Germán Amézaga s/n, Edificio Jorge Basadre, Ciudad Universitaria, Lima, Perú. E-mail: <nestor.allgas@gmail.com>, **Sam Shanee**, ANPC and Neotropical Primate Conservation, 23 Portland Road, Manchester, United Kingdom, M32, **Ana Peralta**, ANPC y Grupo Vida Silvestre, Av. Rinconada s/n, Pachacamac, Lima, Perú, y **Noga Shanee**, ANPC y Neotropical Primate Conservation, 23 Portland Road, Manchester, United Kingdom, M32.

PRIMATAS DA RESERVA PARTICULAR DO PATRIMÔNIO NATURAL ÁGUA BOA, CACOAL, RONDÔNIA, BRASIL

*Almérío Câmara Gusmão
Marcella Alves Crispim
Marcelo Lucian Ferronato
José de Sousa e Silva Júnior*

O conhecimento da fauna de primatas do Estado de Rondônia, o qual está inserido no “arco do desmatamento” (Ferreira et al., 2005), é escasso (Ferrari et al, 1996; van Roosmalen et al., 2002). Segundo van Roosmalen et al. (2002), os fragmentos florestais da região podem abrigar até nove espécies. Neste estudo realizamos um levantamento dos primatas habitantes de um fragmento de floresta de terra firme, a Reserva Particular do Patrimônio Natural Água Boa.