

## A SPECIES LIST FOR THE NEW WORLD PRIMATES (PLATYRRHINI): DISTRIBUTION BY COUNTRY, ENDEMISM, AND CONSERVATION STATUS ACCORDING TO THE MACE-LAND SYSTEM

**Anthony B. Rylands**, Departamento de Zoologia, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, 31270-901 Belo Horizonte, Minas Gerais, and Conservation International do Brasil, Avenida Antônio Abrahão Caram 820/302, 31275-000 Belo Horizonte, Minas Gerais, Brazil, **Russell A. Mittermeier**, Conservation International, 1015 Eighteenth Street N.W., Washington, D. C. 20036, USA, and **Ernesto Rodríguez Luna**, Instituto de Neuroetología, Universidad Veracruzana, Apartado Postal 566, Xalapa, Veracruz 91000, México.

### Introduction

The importance of an up to date evaluation of the diversity of Neotropical primates is emphasized by the increasing numbers of threatened species and subspecies, and the need to pinpoint specific areas for priority action for their protection. The latter may be at the country level (the "megadiversity country" concept: Mittermeier 1987; Mittermeier and Oates 1985; Mittermeier *et al.* 1992b; Conservation International, 1990) or at the level of biomes, regions or localities, for example the "hot-spot" concept of Myers (1988, 1990), and regional workshops to establish priority conservation areas in specific biomes (for example, Fonseca *et al.* 1995).

Priority setting at the species or species' group level has also been emphasized in recent years with the elaboration of action plans by the Specialist Groups of the IUCN Species Survival Commission (Mittermeier, 1987; McNeely *et al.* 1990) along with a major revision of the methodology involved in drawing up the *IUCN Red List of Threatened Animals* (Groombridge, 1994) and the evaluation of their threatened status. The Mace-Lande system for categorizing threatened species was adopted by the World Conservation Union (IUCN) in December 1994 (IUCN 1994), and includes three levels of threat (Critically Endangered, Endangered, and Vulnerable), a further four related categories (Extinct, Extinct in the Wild, Lower Risk, Data Deficient) and a final category (Not Evaluated) which allows for situations where assessment is not possible. As such, the Mace-Lande system requires a working list of *all* species and subspecies of platyrhines. It was first proposed in 1991 (Mace and Lande 1991) and was subsequently amply discussed and revised. Four versions were published before it was finally accepted by IUCN (Mace and Lande 1991; Mace *et al.* 1992; IUCN 1993; Mace and Stuart 1993). The process involved in establishing this new system and the philosophy behind it are discussed in these publications and also Mace (1994a, 1994b, 1995).

Here we present a summary review of the recent taxonomic studies published for Neotropical primates, with the specific aim of establishing a species list for the application of the Mace-Lande categories as well as to provide the basis for analyses of distributions by country and by major phytogeographic region (based on Gentry, 1982) for the evaluation of key areas.

### Taxonomy - Families and Subfamilies

The taxonomy of the New World primates has undergone considerable change over the last two decades, stimulated in large part by the extensive revision of the callitrichids by Professor Philip Hershkovitz (1977, 1979, 1982), and his subsequent reviews of the cebid genera, *Aotus*, *Saimiri*, *Chiropotes*, *Pithecia*, *Cacajao* and *Callicebus* (1983, 1984, 1985, 1987a, 1987b, 1990). Considerable attention has also been given to the phylogeny and taxonomy at family and subfamily level, principally through morphological studies (Rosenberger 1980, 1981; Rosenberger and Coimbra-Filho 1984; Rosenberger and Strier 1989; Rosenberger *et al.* 1990), but also more recently using chromosome and molecular genetics (Dutrillaux 1988; Schneider *et al.* 1993, 1995; Schneider *et al.* in press). Cytotaxonomy is also becoming increasingly important for systematics at the species and subspecies level (see for, example the revision of *Aotus* by Hershkovitz, 1983).

Platyrhine systematics at the family and subfamily level were reviewed by Rosenberger (1981). The predominant classification during this century has involved the use of just two families, the Callitrichidae (or, formerly, Hapalidae) and Cebidae, with *Callimico* being placed in either of the two, or in its own family (v. Dollman 1933; Hershkovitz 1977). This system was maintained in all of the major syntheses published over the last 75 years (for example, Simpson 1945; Hill 1957, 1960, 1962; Cabrera 1957; Napier and Napier 1967; Simons 1972; and Hershkovitz 1977). Hershkovitz (1977) placed the extant cebids in seven

subfamilies: Saimiriinae, Aotinae, Callicebinae, Alouattinae, Pitheciinae, Cebinae, and Atelinae (Table 1). Groves (1989) divided the Cebidae into five families (Cebidae, Aotidae, Atelidae, Callicebidae, and Pitheciidae), although he subsequently (1993) modified this arrangement, defining the Cebidae by the following subfamilies: Alouattinae, Aotinae, Atelinae (*Ateles*, *Brachyteles*, and *Lagothrix*), Callicebinae, Cebinae (including *Cebus* and *Saimiri*, following Rosenberger 1981, see below), and Pitheciinae (*Cacajao*, *Chiropotes* and *Pithecia*).

Rosenberger (1981) proposed an alternative classification (Table 1), based on hypotheses concerning phylogenetic relationships and a proposal for the scenario of their adaptive radiation (see also, Rosenberger, 1980; Rosenberger and Coimbra-Filho, 1984; Rosenberger and Strier, 1989; Rosenberger *et al.* 1990). Rosenberger's (1981) scheme includes the use of tribes and subtribes (Table 1). It maintains but redefines the use of the family name Cebidae to include two subfamilies; the Cebinae (extant genera *Cebus* and *Saimiri*) and Callitrichinae (extant genera *Callithrix* [includes *Cebuella*], *Saguinus*, *Leontopithecus*, and *Callimico*), and places all other platyrhines into a second family, the Atelidae, which includes the following subfamilies: Atelinae (*Alouatta*, *Ateles*, *Brachyteles*, *Lagothrix*) and Pitheciinae (*Pithecia*, *Chiropotes*, *Cacajao*, *Callicebus*, *Aotus*).

A recent molecular phylogeny, elaborated by Schneider *et al.* (1993; 1995, Schneider *et al.* in press), agrees in large part with the classification of Rosenberger (1981), having only one major difference; the alignment of *Aotus* with the Cebidae (subfamily Aotinae) rather than the Pitheciinae. Tyler (1991) argued that there are three-well-defined clades among the extant platyrhines: atelines, pithecines, and callitrichids, but that evolutionary relationships among the remaining four genera (*Aotus*, *Callicebus*, *Cebus* and *Saimiri*) remain uncertain. Like Schneider *et al.* (1993), Tyler (1991) argued that *Aotus* and *Callicebus* are not closely related, but went even further, and, reviewing morphological characters of the former, concluded that "*Aotus* must be removed from other ceboids and made a sister group of the entire Anthropoidea". Dutrillaux (1988) also argued for the primitiveness of *Aotus* on the basis of its karyotype.

The other differences between the schemes of Rosenberger *et al.* (1990) and Schneider *et al.* (1993, 1995, in press) are as follows: 1) Schneider *et al.* place *Saguinus* in the Subtribe Saguina rather than Leontocebina; 2) Schneider *et al.* separate

*Leontopithecus* from *Callithrix* and *Cebuella* and place it in its own Subtribe Leontopithecina; 3) Schneider *et al.* align *Callimico* more closely with *Callithrix* and *Cebuella* and place it in the Subtribe Callimiconina; 4) Schneider *et al.* align *Lagothrix* with *Brachyteles* in the Subtribe Brachytelina, whereas Rosenberger *et al.* place *Lagothrix* in the Subtribe Atelina 5) Schneider *et al.* place *Cacajao* and *Chiropotes* in the Subtribe Chiropotina separate from *Pithecia* and; 5) place *Callicebus* in a separate Tribe rather than Subtribe of the Pitheciinae.

While not making any statement regarding the validity of the various classifications outlined above, the species list we present in the next section maintains the division of Callitrichidae Gray 1821 and Cebidae Bonaparte 1831 as used by Hershkovitz (1977) (Table 1).

### Species and Subspecies

The taxonomy of most of the platyrhine genera is still controversial, with discrepancies arising due to differing emphases given to the concepts and characters used in describing and differentiating forms at the species and subspecies level. Table 2 and Table 3 give a listing of the callitrichid and cebid species and subspecies. We have provided notes so that the reader can be aware of differing opinions, but we emphasize that the lists do not reflect any judgment on our part as to the validity or otherwise of the numerous opposing points of view. We have taken, however, the side of the "splitters" rather than the "lumpers". This is most evident in cases such as that of *Saimiri*, where current evaluations have demonstrated the existence of five species and 12 subspecies on the one hand (Hershkovitz 1984, 1987b), and just two (or even one) species, on the other (Costello *et al.* 1993). *Alouatta* is another case in point, where contemporary taxonomic research is only just emerging. Our aim is to provide an estimate of the diversity of Neotropical primates, and while there is still discussion as to the validity or otherwise of numerous forms, we prefer to maintain them. For conservation purposes it would be a serious mistake to ignore the genetic and morphological diversity of such wide-ranging genera as *Saimiri*. Even while splitting, it is highly probable that the list understates the diversity of most Neotropical primate genera, most particularly in such as *Ateles*, *Alouatta*, *Cebus*, and *Lagothrix*, which have yet to receive modern systematic revisions, but also in, for example *Callithrix*, for which new species have been described recently, and more will undoubtedly be described in the near future.

The listings in Tables 2 and 3 provide a total of five genera, 35 species and 55 species and subspecies of callitrichids, and 11 genera, 63 species and 147 species and subspecies of cebids. In all, we list 16 genera, 98 species and 202 species and subspecies of Neotropical monkeys, ranging from southern Mexico through to northern Argentina and Paraguay, and possibly the northeasternmost tip of Uruguay (Table 9). Mittermeier (1987), in his analysis of the diversity of Neotropical primates and their distributions by country and the major phytogeographic regions of Gentry (1982), worked on the basis of "some 65 species". The increase in species listed here comes from both taxonomic revisions which have raised subspecific forms to full species (for example, the

placing of all *Callithrix* forms as species, and the recognition of such as *Alouatta sara*, *A. arctoidea*, *Ateles chamek*, *Ateles marginatus*, and *Brachyteles hypoxanthus*), as well the descriptions of entirely new species, including *Callithrix mauesi*, *Callithrix marcai*, *Callithrix saterei*, *Leontopithecus caissara*, *Aotus hershkovitzii*, *Callicebus dubius* and *Cebus kaapori*. The taxonomic revision of *Callicebus* by Hershkovitz (1990) increased the number of species from just three to 13, and a further species has yet to be described (Kobayashi and Langguth 1994a, 1994b).

Table 1. The classifications of the extant genera of Platyrrhini according to Hershkovitz (1977), Rosenberger (1981; Rosenberger *et al.* 1990), and Schneider *et al.* (1993, in press).

Hershkovitz (1977)	Rosenberger (1981)	Schneider <i>et al.</i> (in press)
Infraorder Platyrrhini	Infraorder Platyrrhini	Infraorder Platyrrhini
Family Callitrichidae	Family Cebidae	Family Cebidae
<i>Cebuella</i> Gray 1886	Subfamily Cebinae	Subfamily Cebinae
<i>Callithrix</i> Erxleben 1777	Tribe Cebini	Tribe Cebini
<i>Leontopithecus</i> Lesson 1840	<i>Cebus</i> Erxleben 1777	<i>Cebus</i> Erxleben 1777
<i>Saguinus</i> Hoffmannsegg 1807	Tribe Saimiriini	Tribe Saimiriini
Family Callimiconidae	<i>Saimiri</i> Voigt 1831	<i>Saimiri</i> Voigt 1831
<i>Callimico</i> Thomas 1913	Subfamily Callitrichinae	Subfamily Aotinae
Family Cebidae	Tribe Callitrichini	<i>Aotus</i> Illiger 1811
Subfamily Saimiriinae	Subtribe Callitrichina	Subfamily Callitrichinae
<i>Saimiri</i> Voigt 1831	<i>Callithrix</i> Erxleben 1777	Tribe Callitrichini
Subfamily Aotinae	<i>Cebuella</i> Gray 1866	Subtribe Callitrichina
<i>Aotus</i> Illiger 1811	<i>Leontopithecus</i> Lesson 1840	<i>Callithrix</i> Erxleben 1777
Subfamily Callicebinae	Subtribe Leontocebina	<i>Cebuella</i> Gray 1866
<i>Callicebus</i> Thomas 1903	<i>Saguinus</i> Hoffmannsegg 1807	Subtribe Saguina
Subfamily Alouattinae	Tribe Callimiconini	<i>Saguinus</i> Hoffmannsegg 1807
<i>Alouatta</i> Lacépède 1799	<i>Callimico</i> Thomas 1913	Subtribe Leontopithecina
Subfamily Pitheciinae	Family Atelidae	<i>Leontopithecus</i> Lesson 1840
<i>Pithecia</i> Desmarest 1820	Subfamily Atelinae	Subtribe Callimiconina
<i>Chiropotes</i> Lesson 1840	Tribe Atelini	<i>Callimico</i> Thomas 1913
<i>Cacajao</i> Lesson 1840	Subtribe Atelina	Family Atelidae
Subfamily Cebinae	<i>Ateles</i> É. Geoffroy 1806	Subfamily Atelinae
<i>Cebus</i> Erxleben 1777	<i>Brachyteles</i> Spix 1831	Tribe Atelini
Subfamily Atelinae	Subtribe Lagotrichina	Subtribe Atelina
<i>Ateles</i> É. Geoffroy 1806	<i>Lagothrix</i> É. Geoffroy 1812	<i>Ateles</i> É. Geoffroy 1806
<i>Lagothrix</i> É. Geoffroy 1812	Tribe Alouattini	Subtribe Brachytelina
<i>Brachyteles</i> Spix 1831	<i>Alouatta</i> Lacépède 1799	<i>Brachyteles</i> Spix 1831
	Subfamily Pitheciinae	<i>Lagothrix</i> É. Geoffroy 1812
	Tribe Pitheciini	Tribe Alouattini
	Subtribe Pitheciina	<i>Alouatta</i> Lacépède 1799
	<i>Pithecia</i> Desmarest 1820	Subfamily Pitheciinae
	<i>Chiropotes</i> Lesson 1840	Tribe Pitheciini
	<i>Cacajao</i> Lesson 1840	Subtribe Pitheciina
	Subtribe Aotina	<i>Pithecia</i> Desmarest 1820
	<i>Aotus</i> Illiger 1811	Subtribe Chiropotina
	Subtribe Callicebina	<i>Chiropotes</i> Lesson 1840
	<i>Callicebus</i> Thomas 1903	<i>Cacajao</i> Lesson 1840
		Tribe Callicebini
		<i>Callicebus</i> Thomas 1903

Table 2. A listing of the species and subspecies of the Family Callitrichidae.

Family Callitrichidae	Notes	Common name (1)
<i>Cebuella</i> Gray 1866	2	Pygmy marmoset
<i>C. pygmaea</i> Spix 1823		Pygmy marmoset
<i>Callithrix</i> Erxleben 1777	3	Marmosets
<b><i>Callithrix argentata</i> Group</b>		
<i>C. argentata</i> (Linnaeus 1771)	3,4	Bare-ear and tassel-ear marmosets
<i>C. leucippe</i> (Thomas 1922)	3	Silvery marmoset
<i>C. melanura</i> (É. Geoffroy in Humboldt 1812)	3	Golden-white bare-ear marmoset
<i>C. intermedia</i> Hershkovitz 1977	3,4	Black-tailed marmoset
<i>C. emiliae</i> (Thomas 1920)	3,4,5	Aripuanã marmoset
<i>C. nigriceps</i> Ferrari & Lopes 1992	3,6	Snehlage's marmoset
<i>C. marcai</i> Alperin 1993	3,7	Black-headed marmoset
<i>C. humeralifera</i> (É. Geoffroy in Humboldt 1812)	3,4	Marca's marmoset
<i>C. chrysoleuca</i> (Wagner 1842)	3	Black and white tassel-ear marmoset
<i>C. mauesi</i> Mittermeier, Schwarz & Ayres 1992	3,8	Golden-white tassel-ear marmoset
<i>C. saterei</i> Silva e Sousa Jr & Noronha 1995	3,9	Maués marmoset
		Sateré marmoset
<b><i>Callithrix jacchus</i> Group</b>		True marmosets
<i>C. jacchus</i> (Linnaeus 1758)	3	White-tufted-ear marmoset, common marmoset
<i>C. penicillata</i> (É. Geoffroy 1812)	3	Black-tufted-ear marmoset
<i>C. kuhli</i> (Wied-Neuwied 1826)	3,9	Wied's black-tufted-ear marmoset
<i>C. geoffroyi</i> (É. Geoffroy in Humboldt 1812)	3	Geoffroy's tufted-ear marmoset
<i>C. aurita</i> (É. Geoffroy in Humboldt 1812)	3,10	Buffy-tufted-ear marmoset
<i>C. flaviceps</i> (Thomas 1903)	3,11	Buffy-headed marmoset
<i>Saguinus</i> Hoffmannsegg 1807		Tamarins
<b>Hairy-face tamarin Section</b>		
<b><i>Saguinus nigricollis</i> Group</b>		White-mouth tamarins
<i>S. nigricollis nigricollis</i> (Spix 1823)	12	Spix's black mantle tamarin
<i>S. nigricollis graellsii</i> (Jiménez de la Espada 1870)	12	Graell's black-mantle tamarin
<i>S. nigricollis hernandezii</i> Hershkovitz 1982	12	Hernández-Camacho's black mantle tamarin
<i>S. fuscicollis fuscicollis</i> (Spix 1823)	13	Spix's saddle-back tamarin
<i>S. fuscicollis fuscus</i> (Lesson 1840)	13	Lesson's saddle-back tamarin
<i>S. fuscicollis avilapiresi</i> Hershkovitz 1966	13	Ávila Pires' saddle-back tamarin
<i>S. fuscicollis cruzlimai</i> Hershkovitz 1966	13	Cruz Lima's saddle-back tamarin
<i>S. fuscicollis leucogenys</i> (Gray 1866)	13	Andean saddle-back tamarin
<i>S. fuscicollis lagonotus</i> (Jiménez de la Espada 1870)	13	Red-mantle saddle-back tamarin
<i>S. fuscicollis primitivus</i> Hershkovitz 1977	13	Saddle-back tamarin
<i>S. fuscicollis illigeri</i> (Pucheran 1845)	13	Illiger's saddle-back tamarin
<i>S. fuscicollis nigrifrons</i> (I. Geoffroy 1850)	13	Geoffroy's saddle-back tamarin
<i>S. fuscicollis weddelli</i> (Deville 1849)	13	Weddell's saddle-back tamarin
<i>S. fuscicollis melanoleucus</i> (Miranda Ribeiro 1912)	13	White saddle-back tamarin
<i>S. fuscicollis crandalli</i> Hershkovitz 1966	13	Crandall's saddle-back tamarin
<i>S. tripartitus</i> (Milne-Edwards 1878)	14	Golden-mantle saddle-back tamarin
<b><i>Saguinus mystax</i> Group</b>		Moustached tamarins
<i>S. mystax mystax</i> (Spix 1823)	15	Spix's moustached tamarin
<i>S. mystax pileatus</i> (I. Geoffroy & Deville 1848)	15	Red-cap moustached tamarin
<i>S. mystax pluto</i> (Lönnberg 1926)	15	White-rump moustached tamarin
<i>S. labiatus labiatus</i> (É. Geoffroy in Humboldt 1812)	16	Geoffroy's moustached tamarin, red-bellied tamarin
<i>S. labiatus thomasi</i> (Goeldi 1907)	16	Thomas' moustached tamarin
<i>S. imperator imperator</i> (Goeldi 1907)	17	Black-chinned emperor tamarin
<i>S. imperator subgrisescens</i> (Lönnberg 1940)	17	Bearded emperor tamarin
<i>S. midas midas</i> (Linnaeus 1758)	18	Golden-handed tamarin
<i>S. midas niger</i> (É. Geoffroy 1803)	18	Black-handed tamarin
<b>Mottled-face tamarin Section</b>		
<i>S. inustus</i> (Schwarz 1951)	19	Mottled-face tamarin

continued ...

Table 2 (continued)

**Bare-face tamarin Section*****Saguinus bicolor* Group**

<i>S. bicolor bicolor</i> (Spix 1823)	20	Brazilian bare-faced tamarins
<i>S. bicolor martinsi</i> (Thomas 1912)	20	Pied bare-face tamarin
<i>S. bicolor ochraceus</i> Hershkovitz 1966	20	Martin's bare-face tamarin

***Saguinus oedipus* Group**

<i>S. leucopus</i> (Günther 1877)	21	Colombian and Panamanian bare-face tamarins
<i>S. oedipus</i> (Linnaeus 1758)	21	Silvery-brown bare-face tamarin
<i>S. geoffroyi</i> (Pucheran 1845)	21	Cotton-top tamarin

***Leontopithecus* Lesson 1840**

<i>L. rosalia</i> (Linnaeus 1766)	22	Lion tamarins
<i>L. chrysomelas</i> (Kuhl 1820)	22	Golden lion tamarin
<i>L. chrysopygus</i> (Mikan 1823)	22	Golden-headed lion tamarin
<i>L. caissara</i> Lorini & Persson 1990	22	Black lion tamarin
		Black-faced lion tamarin

***Callimico* Miranda Ribeiro 1911**

<i>C. goeldii</i> (Thomas 1904)	23	Goeldi's monkey
---------------------------------	----	-----------------

**Notes**

1. Common names of Callitrichidae follow Hershkovitz (1977).

2. Rosenberger (1981 see also Rosenberger and Coimbra-Filho, 1984) argued, on morphological terms, that the pygmy marmoset should correctly be included in the genus *Callithrix*. This argument was not maintained in Rosenberger *et al.* (1990), although recently Barroso (1995) and Schneider *et al.* (in press) also argued for this arrangement on the basis of molecular genetics. Taxonomy and distribution follows Hershkovitz (1977), Eisenberg (1989) and Bicca-Marques and Calegaro-Marques (1995).

3. Hershkovitz (1977) placed all of the *Callithrix jacchus* Group marmosets as subspecies of *Callithrix jacchus*. Coimbra-Filho (1970a, 1971, 1990) and Coimbra-Filho and Mittermeier (1973b), Mittermeier and Coimbra-Filho (1981), Natori (1986, 1994), Vivo (1985, 1991), Mittermeier *et al.* (1988, 1992a), Natori and Shigehara (1992), and Groves (1993) (see also Marroig, 1995) listed all the Atlantic forest marmosets as full species. Coimbra-Filho (1990) argued that the three subspecies of *C. humeralifer* (*C. h. humeralifer*, *C. h. chrysoleuca*, and *C. h. intermedius*) listed by Hershkovitz (1977) should be considered species, following De Boer (1974) and Vivo (1991). Genetic studies by Meireles *et al.* (1992) concluded that *C. humeralifer* and *C. emiliae* are subspecies of *C. argentata*; and that *C. jacchus*, *C. penicillata*, and *C. geoffroyi* are subspecies; but that *C. geoffroyi* is the most primitive of the "jacchus" group (most similar genetically to *C. argentata*), which should therefore be referred to as the "geoffroyi" group, with *C. jacchus* and *C. penicillata* as subspecific to *C. geoffroyi*. The primitiveness (Hershkovitz 1977) of *C. geoffroyi* was also advocated by Natori (1986, 1994) and Natori and Shigehara (1992), and also makes sense concerning the degree of morphological adaptation for tree-gouging, which is most advanced in *C. jacchus* and *C. penicillata*. The following publications list

*all* marmosets as species: Mittermeier *et al.* (1992a), Vivo (1985, 1991), and Groves (1993). Distributions of *Callithrix* according to Hershkovitz (1977), Stallings (1985), Mittermeier *et al.* (1992a), and Rylands *et al.* (1993).

4. *Callithrix intermedia* was described as an intermediate color form of *C. humeralifer* by Hershkovitz (1977). Pelage color and pattern, and its geographic distribution, however, would place it as a subspecies of *C. argentata* if, following Hershkovitz (1977), the Amazonian marmosets are considered subspecies of just two species, *C. argentata* and *C. humeralifer* (see Rylands *et al.* 1993). Meireles *et al.* (1992) indicated that *C. humeralifer* and *C. emiliae* are subspecific to *C. argentata*.

5. *Callithrix emiliae* was first described by Thomas (1920) from the Rio Iriri, southern Pará. It was not recognized by Hershkovitz (1977) who regarded it as a dark form of *C. argentata* *argentata*. Vivo (1985; see also Vivo 1991), revalidated this form on the basis of specimens from the state of Rondônia. However, Rylands *et al.* (1993) argued that the Rondônia marmosets described by Vivo should be considered a distinct species based on the fact that the distribution of this form and that of the *C. emiliae* described by Thomas (1920) from Maloca on the Rio Curuá (see Vivo 1985; Ávila Pires 1986) are disjunct, and separated by *C. melanura*. *C. emiliae* was not listed by Groves (1993). Distribution according to Vivo (1985, 1991).

6. Distribution of *Callithrix nigriceps* based on Ferrari and Lopes (1992) and Ferrari (1993, 1994).

7. *Callithrix marcai* was described as a subspecies of *Callithrix argentata* by Alperin 1993. It is listed here as a species to conform with the view that all *Callithrix* should be considered species (Vivo 1985, 1991; Mittermeier *et al.* 1992a). It is known only from the type locality, "Foz do Rio Castanho (= Rio Roosevelt), afluente e esquerda do rio Aripuanã. Estado do Amazonas, Brasil" (Alperin 1993). The type locality as described by Alperin is confused in

that the Rio Castanho is not a synonym of the Rio Roosevelt, and is a left bank affluent of the Rio Roosevelt, not the Rio Aripuanã. Confused also is the map and gazetteer of localities provided in Alperin (1993) which do not correspond.

8. Distribution of *Callithrix maeusi* according to Mittermeier *et al.* (1992a) and Silva Jr. and Noronha (1995a).

9. Coimbra-Filho (1984, 1985, 1990), Rylands (1989b), Mittermeier *et al.* (1988), Natori (1990), Rylands *et al.* (1993), and Groves (1993) recognize *Callithrix kuhli* from southern Bahia as a valid form. Hershkovitz (1977) regarded it as a hybrid between *C. j. geoffroyi* and *C. j. penicillata*. Vivo (1991) considered it to be indistinguishable from *C. penicillata*. Distribution restricted to Brazil (southern Bahia) follows Rylands *et al.* (1993). Groves (1993) gave the distribution as Bolivia - evidently a typographic error.

10. The distribution of *Callithrix aurita* is reviewed by Hershkovitz (1977), Muskin (1984), Coimbra-Filho (1986b, 1991), and Olmos and Martuscelli (1995).

11. Coimbra-Filho (1986a, 1986b, 1990; Coimbra-Filho *et al.* 1991) argued that *Callithrix flaviceps* should be considered subspecific to *C. aurita*. The distribution of *C. flaviceps* is reviewed by Hershkovitz (1977), Coimbra-Filho (1986a), and Mendes (1993).

12. The taxonomy of *Saguinus nigricollis* follows Hershkovitz (1982). *S. n. graellsii* is listed as a full species, *S. graellsii*, by Hernández-Camacho and Cooper (1976) and Defler (1994), on the basis that it is sympatric with a population of *S. nigricollis* in the region of Puerto Leguízamo in southern Colombia. Hernández-Camacho and Defler (1991) listed it as a subspecies of *S. nigricollis*, in conformity with Hershkovitz (1977). Defler (1994) stated that further studies are needed to prove the species' status of this form. Distribution according to Hernández-Camacho and Cooper (1976), Hershkovitz (1982) and Eisenberg (1989).

13. The taxonomy of *Saguinus fuscicollis* is based on Hershkovitz (1977; see also Cheverud and Moore 1990). *S. f. melanoleucus*, *S. f. acrensis*, and *S. f. crandalli* were listed as subspecies of *S. melanoleucus* by Coimbra-Filho (1990). *Saguinus fuscicollis acrensis* Carvalho 1957 is not considered a valid form, following Peres (1991). *S. f. cruzlimai* and *S. f. crandalli* are of unknown provenance (Hershkovitz 1977). Hernández-Camacho and Cooper (1976) indicated the existence of an undescribed *S. fuscicollis* subspecies in the region of San José de Guaviare, Colombia (also cited by Defler 1994). Distributions are given by Hershkovitz (1977), Eisenberg (1989), and Aquino and Encarnación (1995) (see also Rylands *et al.* 1993).

14. Hershkovitz (1977) listed *Saguinus tripartitus* as a subspecies of *S. fuscicollis*. Thorington (1988) argued for its species status (see also Albuja, 1994;). Distribution based on Hershkovitz (1977), Thorington (1988), Albuja (1994), and Aquino and Encarnación (1994, in press).

15. The taxonomy of *Saguinus mystax* follows Hershkovitz (1977). Distributions are based on Hershkovitz (1977), Rylands *et al.* (1993), and Aquino and Encarnación (1994).

16. The taxonomy of *Saguinus labiatus* follows Hershkovitz (1977). Distribution based on Hershkovitz (1977) and Silva Jr. (1988), Aquino and Castro (1989), and Aquino and Encarnación (1994).

17. The taxonomy of *Saguinus imperator* follows Hershkovitz (1982). Distribution based on Hershkovitz (1982), Aquino and Encarnación (1994).

18. The taxonomy and distribution of *Saguinus midas* follows Hershkovitz (1977) and Eisenberg (1989). On the basis of morphometric studies of the postcanine dentition, Natori and Hanihara (1992) found *S. m. midas* to be more similar to *S. bicolor* than to *S. m. niger*. For this reason *S. m. niger* should possibly be raised to species status (Rylands *et al.* 1993), and would also argue for a modification of the species' group arrangements of Hershkovitz (1977): that is, placing the *S. midas* Group in the Bare-face Tamarin section. Melo *et al.* (1992), on the other hand, examined blood genetic systems in the two subspecies and obtained results compatible with their classification as subspecies, not being sufficiently divergent to warrant species status.

19. The taxonomy and distribution of *Saguinus inustus* follows Hernández-Camacho and Cooper (1976), Hershkovitz (1977), Eisenberg (1989), and Rylands *et al.* (1993). Hernández-Camacho and Defler (1991) indicated possibility of two subspecies in Colombia.

20. The taxonomy of *Saguinus bicolor* is based on Hershkovitz (1977). Distributions follow Hershkovitz (1977), Ayres *et al.* (1982), Egler (1983) and Coimbra-Filho (1987).

21. Hershkovitz (1977) considered *Saguinus geoffroyi* to be subspecific to *S. oedipus*. Comparative morphological studies by Hanihara and Natori (1987), Moore and Cheverud (1992) and Skinner (1991) argue for them being separate species (see also Rylands, 1993). Hernández-Camacho and Cooper (1976), Mittermeier and Coimbra-Filho (1981), Hernández-Camacho and Defler (1985, 1991), Mittermeier *et al.* (1988), Rylands *et al.* (1993), Groves (1993), Mast *et al.* (1993) and Defler (1994) considered them to be separate species. Distributions are based on Hernández-Camacho and Cooper (1976), Hershkovitz (1977), Eisenberg (1989), Hernández-Camacho and Defler (1991), and Mast *et al.* (1993). Eisenberg (1989) lists *S. geoffroyi* and *S. oedipus* as separate species. The author of the name *S. geoffroyi* is given by him as Reichenbach 1862.

22. The lion tamarins, *Leontopithecus*, are given as separate species following Della Serra (1951), Rosenberger and Coimbra-Filho (1984), Mittermeier *et al.* (1988), Natori (1989), and Rylands *et al.* (1993). They have been listed as subspecies of *L. rosalia* by Coimbra-Filho and Mittermeier (1972, 1973a), Hershkovitz (1977), Mittermeier and Coimbra-Filho (1981), and Forman *et al.* (1986). *L. caissara*

Lorini and Persson 1990 is regarded as a subspecies of *L. chrysopygus* by Coimbra-Filho (1990). Distribution follows Coimbra-Filho and Mittermeier (1972, 1973a, 1977), Hershkovitz (1977), Rylands *et al.* (1991), Rosenberger and Coimbra-Filho (1984), Kierulff (1993), Kierulff and Oliveira (1994), Pinto and Tavares (1994), Lorini and Persson (1994), and Valladares-Padua *et al.* (1994a, 1994b).

23. Hershkovitz (1977) placed *Callimico* in its own family, Callimiconidae. Rosenberger (1981) argued for its alignment as the subfamily Callimiconinae within the family Cebidae, redefined as including, besides, the subfamilies Cebinae (*Cebus, Saimiri*) and Callitrichinae. See also Martin (1991) and Barroso (1995). Distribution according to Hershkovitz (1977) and Eisenberg (1989).

Table 3. A listing of the species and subspecies of the Family Cebidae.

Family Cebidae	Notes	Common name
<i>Aotus</i> Illiger 1811	1-2	Night monkeys, owl monkeys, douroucouli
<b>Gray-neck Species Group</b>		
<i>A. lemurinus lemurinus</i> (I. Geoffroy 1846)	1	Colombian or lemuring night monkey
<i>A. lemurinus griseimembra</i> (Elliot 1913)	1	Grey-legged night monkey
<i>A. vociferans</i> (Spix 1823)		
<i>A. trivirgatus</i> (Humboldt 1812)	1	Douroucouli, owl monkey, night monkey
<i>A. brumbacki</i> Hershkovitz 1983	1	Brumback's night monkey
<i>A. hershkovitzii</i> Ramirez-Cerquera 1983	1	Hershkovitz's night monkey
<b>Red-neck Species Group</b>		
<i>A. miconax</i> (Thomas 1927)	1	Andean night monkey
<i>A. nigriceps</i> Dollman 1909	1	Black-headed or Peruvian night monkey
<i>A. infulatus</i> (Kuhl 1820)	1	Feline night monkey
<i>A. azarae azarae</i> (Humboldt 1812)	1,2	Azara's night monkey
<i>A. azarae boliviensis</i> (Elliot 1907)	1,2	Bolivian night monkey
<i>A. nancymaae</i> Hershkovitz 1983	1,2	Ma's night monkey
<i>Callicebus</i> Thomas 1903	3-4	Titi monkeys
<b>Callicebus modestus Group</b>		
<i>C. modestus</i> Lönnberg 1939	3	
<b>Callicebus donacophilus Group</b>		
<i>C. donacophilus donacophilus</i> D'Orbigny 1836	3	
<i>C. donacophilus pallescens</i> Thomas 1907	3	
<i>C. olallae</i> (Lönnberg 1939)	3	Beni titi monkey
<i>C. oenanthe</i> (Thomas 1924)	3	Andean titi monkey
<b>Callicebus moloch Group</b>		
<i>C. cinerascens</i> (Spix 1823)	3	
<i>C. hoffmannsi hoffmannsi</i> (Thomas 1908)	3	Hoffmann's titi monkey
<i>C. hoffmannsi baptista</i> (Lönnberg 1939)	3	
<i>C. moloch</i> Hoffmannsegg 1807	3	
<i>C. brunneus</i> (Wagner 1842)	3	
<i>C. cupreus cupreus</i> (Spix 1823)	3	
<i>C. cupreus discolor</i> (I. Geoffroy & Deville 1848)	3	
<i>C. cupreus ornatus</i> (Gray 1866)	3	Ornate titi monkey
<i>C. caligatus</i> (Wagner 1842)	3	
<i>C. dubius</i> Hershkovitz 1988	3	
<i>C. personatus personatus</i> (E. Geoffroy 1812)	3	Northern masked titi
<i>C. personatus nigrifrons</i> (Spix 1823)	3	Black-fronted titi
<i>C. personatus melanochir</i> (Wied-Neuwied 1820)	3	Southern Bahian masked titi
<i>C. personatus barbarabrowniae</i> Hershkovitz 1990	3	Northern Bahian blond titi
<i>Callicebus</i> sp.	3,4	
<b>Callicebus torquatus Group</b>		
<i>C. torquatus torquatus</i> (Hoffmannsegg 1807)	3	Collared titi, widow monkey
<i>C. torquatus medemi</i> Hershkovitz 1963	3	
<i>C. torquatus lugens</i> (Humboldt 1811)	3	
<i>C. torquatus lucifer</i> (Thomas 1914)	3	
<i>C. torquatus purinus</i> (Thomas 1927)	3	
<i>C. torquatus regulus</i> Thomas 1927	3	

continued ...

Table 3 (continued)

<b>Saimiri Voigt 1831</b>	5	Squirrel monkeys
<b>Saimiri boliviensis Group (Roman type)</b>		
<i>S. boliviensis boliviensis</i> (I. Geoffroy & de Blainville 1834)	5	Black-headed squirrel monkey
<i>S. boliviensis peruviensis</i> Hershkovitz 1984	5	Peruvian squirrel monkey
<i>S. boliviensis pluvialis</i> Lönnberg 1940	5	
<i>S. boliviensis jaburuensis</i> Lönnberg 1940	5	
<i>S. vanzolinii</i> Ayres 1981	5	
<b>Saimiri sciureus Group (Gothic type)</b>		
<i>S. sciureus sciureus</i> (Linnaeus 1758)	5	Common squirrel monkey
<i>S. sciureus macrodon</i> (Elliot 1907)	5	Ecuadorian squirrel monkey
<i>S. sciureus cassiquiarensis</i> (Lesson 1840)	5	Humboldt's squirrel monkey
<i>S. sciureus albigena</i> (von Pusch 1941)	5	
<i>S. oerstedi oerstedi</i> (Reinhardt 1872)	5	Black-crowned Central American squirrel monkey
<i>S. oerstedi citrinellus</i> Thomas 1904	5	Grey-crowned Central American squirrel monkey
<i>S. ustus</i> I. Geoffroy 1843	5	Golden-backed squirrel monkey
<b>Cebus Erxleben 1777</b>	6-7	Tufted and untufted capuchins
<b>Tufted Group</b>		
<i>C. apella apella</i> (Linnaeus 1758)	6	Guianan brown capuchin
<i>C. apella macrocephalus</i> Spix 1823	6	Large-headed capuchin
<i>C. apella maranensis</i> (von Pusch, 1941)	6	Marañón tufted capuchin
<i>C. apella pallidus</i> (Gray 1865)	6	Tambopata tufted capuchin, pale capuchin
<i>C. apella peruanus</i> (Thomas 1901)	6	Peruvian tufted capuchin
<i>C. apella libidinosus</i> (Spix 1823)	6	Bearded capuchin
<i>C. apella nigritus</i> (Goldfuss 1809)	6	Black-horned capuchin
<i>C. apella robustus</i> (Kuhl 1820)	6	Crested capuchin
<i>C. apella margaritae</i> Hollister 1914	6	Margarita Island capuchin
<i>C. apella paraguayanus</i> Fischer, 1829	6	Paraguayan tufted capuchin
<i>C. xanthosternos</i> (Wied 1820)	6,7	Yellow-breasted capuchin
<b>Untufted Group</b>		
<i>C. albifrons albifrons</i> (Humboldt 1812)	8	White-fronted capuchin
<i>C. albifrons unicolor</i> Spix 1823	8	
<i>C. albifrons adustus</i> Hershkovitz 1949	8	Brown-faced capuchin
<i>C. albifrons aequatorialis</i> Allen 1914	8	Ecuadorian capuchin
<i>C. albifrons cesareae</i> Hershkovitz 1949	8	
<i>C. albifrons cuscinus</i> (Thomas 1901)	8	Shock-headed capuchin
<i>C. albifrons malitiosus</i> Elliot 1909	8	
<i>C. albifrons trinitatis</i> von Pusch 1941	8	Trinidad white-fronted capuchin
<i>C. albifrons versicolor</i> Pucheran 1845	8	Varied capuchin
<i>C. albifrons yuracus</i> Hershkovitz 1949	8	Andean white-fronted capuchin
<i>C. albifrons leucocephalus</i> Gray 1865	8	
<i>C. capucinus capucinus</i> (Linnaeus 1758)	9	White-throat capuchin, white-faced capuchin
<i>C. capucinus limitaneus</i> Hollister 1914	9	
<i>C. capucinus imitator</i> Thomas 1903	9	Panamanian white-throated capuchin
<i>C. capucinus curtus</i> Bangs 1905	9	Gorgona white-fronted capuchin
<i>C. olivaceus olivaceus</i> Schomburgk 1848	10	Wedge-capped capuchin
<i>C. olivaceus apiculatus</i> Hershkovitz 1949	10	
<i>C. olivaceus brunneus</i> Allen 1914	10	Brown weeper capuchin
<i>C. olivaceus castaneus</i> I. Geoffroy 1851	10	Chestnut capuchin
<i>C. olivaceus</i> ssp.	10,11	
<i>C. kaapori</i> Queiroz 1992	10	Ka'apor capuchin
<b>Pithecia Desmarest 1804</b>	12	Saki monkeys
<i>P. pithecia pithecia</i> Linnaeus 1766	12	White-faced saki
<i>P. pithecia chrysoccephala</i> I. Geoffroy 1850	12	Golden-faced saki
<i>P. monachus monachus</i> (É. Geoffroy 1812)	12	Geoffroy's monk saki
<i>P. monachus milleri</i> Allen 1914	12	Miller's monk saki
<i>P. irrorata irrorata</i> Gray 1842	12	Gray's bald faced saki
<i>P. irrorata vanzolinii</i> Hershkovitz 1987	12	Vanzolini's bald-faced saki
<i>P. albicans</i> Gray 1860	12	White saki, buffy saki

continued ...

Table 3 (continued)

<i>P. aequatorialis</i> Hershkovitz 1987	12	Equatorial saki
<i>Chiropotes</i> Lesson 1840	13	Bearded sakis
<i>C. albinasus</i> (I. Geoffroy & Deville 1848)	13	White-nosed bearded saki
<i>C. satanas satanas</i> (Hoffmannsegg 1807)	13	Bearded saki
<i>C. satanas chiropotes</i> (Humboldt 1811)	13	Black saki
<i>C. satanas utahicki</i> Hershkovitz 1985	13	Uta Hick's bearded saki
<i>Cacajao</i> Lesson 1840	14	Uacaris
<i>C. calvus calvus</i> (I. Geoffroy 1847)	14	White bald-headed uacari
<i>C. calvus ucayalii</i> (Thomas 1928)	14	Ucayali bald-headed uacari
<i>C. calvus novaesi</i> Hershkovitz 1987	14	Novaes' bald-headed uacari
<i>C. calvus rubicundus</i> (I. Geoffroy and Deville 1848)	14	Red bald-headed uacari
<i>C. melanocephalus melanocephalus</i> (Humboldt 1811)	14	Humboldt's black-headed uacari
<i>C. melanocephalus ouakary</i> (Spix 1823)	14	Spix's black-headed uacari
<i>Alouatta</i> Lacépède 1799	15-21	Howling monkeys
<i>A. seniculus seniculus</i> (Linnaeus 1766)	15,21	Red howling monkey
<i>A. seniculus insularis</i> Elliot 1910	15,21	Trinidad howling monkey
<i>A. seniculus stramineus</i> (Humboldt 1812)	15,21	Golden howling monkey
<i>A. seniculus macconnelli</i> Elliot 1910	15,21	Guianan red howling monkey
<i>A. seniculus amazonica</i> Lönnberg 1941	15,21	
<i>A. seniculus juara</i> Elliot 1910	15,21	
<i>A. seniculus puruensis</i> Lönnberg 1941	15,21	
<i>A. seniculus</i> ssp.	16	
<i>A. arctoidea</i> Cabrera 1940	15,21	Ursine howling monkey
<i>A. sara</i> Elliot 1910	15,21	Bolivian red howling monkey
<i>A. belzebul belzebul</i> (Linnaeus 1766)	17,21	Red-handed howling monkey
<i>A. belzebul nigerrima</i> Lönnberg 1941	17,21	Black howling monkey
<i>A. belzebul discolor</i> (Spix 1823)	17,21	
<i>A. belzebul ululata</i> (Elliot 1912)	17,21	
<i>A. fusca fusca</i> (Ihering 1914)	17,21	Northern brown howling monkey
<i>A. fusca clamitans</i> (Cabrera 1940)	17,21	Southern brown howling monkey
<i>A. palliata palliata</i> (Gray 1848)	21	Golden-mantled howling monkey
<i>A. palliata mexicana</i> (Merriam 1902)	21	Mexican howling monkey
<i>A. palliata aequatorialis</i> (Festa 1903)	21	South Pacific blackish howling monkey
<i>A. coibensis coibensis</i> (Thomas 1902)	21	Coiba Island howling monkey
<i>A. coibensis trabeata</i> Lawrence 1933	21	Azuero howling monkey
<i>A. caraya</i> (Humboldt 1812)	19,21	Black howling monkey
<i>A. pigra</i> (Lawrence 1933)	20,21	Black howling monkey
<i>Ateles</i> É. Geoffroy 1806	22-23	Spider monkeys
<i>A. geoffroyi geoffroyi</i> Kuhl 1820	22	Geoffroy's spider monkey
<i>A. geoffroyi azuerensis</i> (Bole 1937)	22	Azuero spider monkey
<i>A. geoffroyi frontatus</i> (Gray 1842)	22	Black-browed spider monkey
<i>A. geoffroyi grisescens</i> (Gray 1866)	22	Hooded spider monkey
<i>A. geoffroyi pan</i> (Schlegel 1876)	22	Guatemalan spider monkey
<i>A. geoffroyi panamensis</i> Kellogg & Goldman 1944	22	Red spider monkey
<i>A. geoffroyi ornatus</i> (Gray 1870)	22	Ornate spider monkey
<i>A. geoffroyi vellerosus</i> (Gray 1866)	22	Mexican spider monkey
<i>A. geoffroyi yucatanensis</i> Kellogg & Goldman 1944	22	Yucatán spider monkey
<i>A. fusciceps fusciceps</i> Gray 1866	22	Brown-headed spider monkey
<i>A. fusciceps robustus</i> (= <i>rufiventris</i> ) (Allen 1914)	22	Colombian black spider monkey
<i>A. chamek</i> (Humboldt 1812)	22	Black-faced black spider monkey
<i>A. paniscus</i> (Linnaeus, 1758)	22	Red-faced black spider monkey
<i>A. marginatus</i> (É. Geoffroy 1809)	22	White-whiskered spider monkey
<i>A. belzebuth belzebuth</i> (É. Geoffroy 1806)	22	White-bellied spider monkey
<i>A. belzebuth hybridus</i> (I. Geoffroy 1829)	22	Variegated spider monkey
<i>A. belzebuth brunneus</i> Gray 1870	22,23	Brown spider monkey

continued ...

Table 3 (continued)

<i>Lagothrix</i> É. Geoffroy 1812	24	Woolly monkeys
<i>L. lagotricha lagotricha</i> (Humboldt 1812)	24	Humboldt's woolly monkey
<i>L. lagotricha cana</i> (É. Geoffroy in Humboldt 1812)	24	Geoffroy's woolly monkey
<i>L. lagotricha poeppigii</i> (Schinz 1844)	24	Poeppig's woolly monkey
<i>L. lagotricha lugens</i> (Elliot 1907)	24	Colombian woolly monkey
<i>L. flavicauda</i> (Humboldt 1812)	24	Yellow-tailed woolly monkey
<i>Brachyteles</i> Spix 1823	25	Muriqui
<i>B. arachnoides</i> (É. Geoffroy 1806)	25	Southern muriqui
<i>B. hypoxanthus</i> (Kuhl 1820)	25	Northern muriqui

## Notes

1. The taxonomy of *Aotus* follows Hershkovitz (1983), except for *A. hershkovitzi*, described by Ramirez-Cerquera in 1983 (*Abstracts. IX Cong. Latinoamericano de Zoología*, Arequipa, Peru, p.148) (see also Defler 1994). Defler (1994) mentions the possibility of a further two Colombian species of *Aotus* from the Colombian trapezium, *A. nigriceps* (possibly introduced) and *A. nancymaae*. Hernández-Camacho and Cooper (1976) referred to *A. trivirgatus zonalis* (Goldman 1914) in Panama and Colombia, which is considered a synonym of *A. t. griseimembra* by Hershkovitz (1949), but as a synonym of *A. l. lemurinus* by Hershkovitz (1983), and a synonym of *A. lemurinus* by Groves (1993). The status recommended by Hershkovitz (1983) was accepted by Hernández-Camacho and Defler (1991). Groves (1993) lists *A. azarai boliviensis* as a synonym of *A. azarai*, and *A. lemurinus griseimembra* as a synonym of *A. lemurinus*. Pieczarka and Nagamachi (1988) found that the karyotypes of *A. infulatus* and *A. azarae boliviensis* are closer to each other than to the karyotype of *A. a. azarae*, and argue for the need to further revise the systematic classification of the genus. Recent cytogenetic studies by Pieczarka *et al.* (1992) have reinforced the validity of the species *A. vociferans* and *A. nancymaae*. A phylogeny based on karyotypes is provided by Galbreath (1983). Distributions according to Hernandez-Camacho and Cooper (1976), Hershkovitz (1983), Stallings (1985), Eisenberg (1989), Zunino *et al.* (1991), Redford and Eisenberg (1992), Fernandes (1993), and Aquino and Encarnación (1994). Common names follow Hill (1960) except in the use of "night monkey" rather than "douroucouli".

2. *A. azarai* is used rather than *A. azarae*, and *A. nancymaae* rather than *A. nancymai* (see Hershkovitz 1983), following Groves (1993) who pointed out the correct specific names in accordance with Article 31 (a(ii)) of the *International Code of Zoological Nomenclature* (International Commission on Zoological Nomenclature, 1985).

3. *Callicebus* taxonomy according to Hershkovitz (1990). Groves (1992, 1993) indicated that *C. brunneus* and *C. hoffmannsi* may be subspecies of *C. moloch* (*C. h. baptista* is a synonym of *C. hoffmannsi*), that *C. caligatus* and *C. dubius* may be synonyms of *C. cupreus*, that *C. cupreus discolor* and *C. c. ornatus* are synonyms of *C. cupreus*, that *C. donacophilus pallescens* is a synonym of *C.*

*donacophilus*, and that all subspecies of *C. torquatus* and *C. personatus* listed here are synonyms of their respective species. Hershkovitz (1990) placed the *Callicebus* species into four groupings: 1) *C. modestus* (*C. modestus*); 2) *C. donacophilus* (*C. olallae*, *C. donacophilus*, *C. oenanthe*); 3) *C. moloch* (*C. brunneus*, *C. caligatus*, *C. cupreus*, *C. dubius*, *C. hoffmannsi*, *C. moloch*, *C. cinerascens* and *C. personatus*); and 4) *C. torquatus* (*C. torquatus*). The species groupings used here are those of Hershkovitz (1990), although a recent phylogenetic study, based on cranial measurements, by Kobayashi (1995) resulted in the following five groupings: 1) *C. donacophilus* (*C. modestus*, *C. olallae*, *C. donacophilus*); 2) *C. cupreus* (*C. caligatus*, *C. cupreus*); *C. moloch* (*C. brunneus*, *C. hoffmannsi*, *C. moloch*, *C. cinerascens*); 4) *C. personatus* (*C. personatus*); and 5) *C. torquatus* (*C. torquatus*). Kobayashi (1995) did not evaluate *C. oenanthe* and *C. p. barbarabrowniae*, and the position of *C. dubius* was uncertain. Kobayashi (1995) indicated that *C. caligatus* is a synonym of *C. cupreus*. Distributions according to Hernández-Camacho and Cooper (1976), Eisenberg (1989), Stallings (1985), Hershkovitz (1990), and Aquino and Encarnación (1994).

4. Kobayashi and Langguth (1994a, 1994b) recorded the existence of a new *Callicebus* from the coast of the state of Sergipe, Brazil.

5. *Saimiri* taxonomy follows Hershkovitz (1984). Hershkovitz (1987b; footnote page 22) also recognized *Saimiri boliviensis jaburuensis* and *S. b. pluvialis* (given as junior synonyms of *S. b. boliviensis* in Hershkovitz 1984) as valid subspecies. Hershkovitz (1987b) referred to *S. vanzolinii* Ayres 1981 as a subspecies of *S. boliviensis*. An alternative taxonomy is presented by Thorington (1985) as follows: *S. sciureus sciureus* (includes the forms *albigena*, *macrodon*, and *ustus* recognized by Hershkovitz, 1984), *S. sciureus boliviensis* (includes the forms *pluvialis* Lönnberg 1940 and *jaburuensis* Lönnberg 1940 recognized by Hershkovitz 1987), *S. sciureus cassiquiarensis*, *S. sciureus oerstedii* (includes the form *citrinellus* recognized by Hershkovitz 1984), and *S. madeirae* (given as a junior synonym of *S. ustus* by Hershkovitz 1984). Hernández-Camacho and Defler (1991) recognize *S. sciureus caquetensis* Allen 1916, given as a junior synonym of *S. sciureus macrodon* by Hershkovitz (1984). Groves (1993) recognized just five species of *Saimiri* (*S. boliviensis*, *S. oerstedii*, *S. sciureus*, *S. ustus* and *S. vanzolinii*) with no subspecies. He listed *S. b. peruviensis*, *S. b. jaburuensis*

and *S. b. pluvialis* as synonyms of *S. boliviensis*, *S. o. citrinellus* as a synonym of *S. oerstedii*, all subspecies of *S. sciureus* listed here as synonyms, and *S. madeirae* recognized by Thorington (1985) as a synonym of *S. ustus*. The most recent review (morphology, genetics and behavior) of squirrel monkey taxonomy by Costello *et al.* (1993) argued for the recognition of just two species: *S. sciureus* in South America, and *S. oerstedii* in Panama and Costa Rica. Silva *et al.* (1993) also gave evidence for just one large polytypic South American species, *Saimiri sciureus*. Distributions according to Hernández-Camacho and Cooper (1976), Hershkovitz (1984), Ayres (1985), Eisenberg (1989), and Aquino and Encarnación (1994).

6. The *C. apella* taxonomy of Hill (1960) was based on personal communications from Kellogg, who was revising the taxonomy but died before completing it. Apart from those listed above, Hill (1960) gives the following subspecies: *Cebus apella fatuellus* Colombian brown capuchin, from northern Colombia, type locality unknown; *Cebus apella tocantinus* from type locality Rio Tocantins - no clear difference from *peruanus*; *Cebus apella magnus* from the Río Putumayo; *Cebus apella juruanus* from the Rio Juruá; *Cebus apella cay* Azara's capuchin from Paraguay, Goiás; *Cebus apella frontatus* the white-cheeked capuchin, no type locality, but argued by Vieira (1955) to be from the Serra da Paranapiacaba, Alto da Serra, Cananéia, Santa Catarina, and Rio Grande do Sul; *Cebus apella vellerosus* from Argentina is given by Hill (1960) as a synonym of *C. a. nigrinus*. Hernández-Camacho and Cooper (1976) argue that *C. apella* north of the Amazon in Colombia exhibit no phenotypic distinctions sufficient to warrant subspecies. They give the subspecies as *C. a. apella* (see also Hernández-Camacho and Defler 1991; Defler 1994). Hershkovitz in his current studies of *Cebus* taxonomy will maintain as valid the subspecies *C. a. maranonis*, *C. a. macrocephalus*, *C. a. pallidus*, and *C. a. peruanus* (*apud* Encarnación, 1993). All these subspecies were listed by Aquino and Encarnación (1994). Groves (1993) gave all subspecific forms listed here as synonyms of *C. apella*. Torres de Assumpção (1988) and Torres (1989) carried out an incomplete study, but the taxonomy of *Cebus apella* awaits a modern revision, and is not definite. Taxonomy of Brazilian Atlantic forest *C. apella* follows Kinzey (1982), except for *C. a. paraguayanus* Fischer, 1829 from Argentina and Paraguay which was considered by him to be a junior synonym of *C. a. libidinosus*. Bodini and Pérez-Hernández (1987) and Bodini (1989) list only two subspecies for Venezuela: *C. a. margaritae* and *C. a. apella* (Federal Territory of Amazonas) (see also Sanz and Márquez, 1994).

7. *C. xanthosternos*, formerly considered a subspecies of *Cebus apella*, is genetically quite distinct (Seuáez *et al.* 1986), and is, therefore, listed as a valid species (Mittermeier *et al.* 1988; see also Coimbra-Filho *et al.* 1991).

8. Eleven subspecies are listed for *Cebus albifrons*. The taxonomy and distributions of the white-fronted capuchins are confused and require revision. The subspecies listed here are based on Hershkovitz (1949) and Hernández-Camacho and Cooper (1976) (see also Hernández-Camacho

and Defler 1985, 1991). Hernández-Camacho and Defler (1991) mentioned six subspecies in Colombia and listed five: *C. a. malitiosus*, *C. a. cesarae*, *C. a. versicolor*, *C. a. albifrons*, and *C. a. yuracus*. They regarded *C. a. unicolor* as possibly a junior synonym of *C. a. albifrons*, although, later, Defler (1994) also included this subspecies for Colombia. Defler (1994) listed six subspecies for Colombia, three endemic: *C. a. albifrons*, *C. a. cesarae* (endemic), *C. a. versicolor* (endemic), *C. a. malitiosus* (endemic), *C. a. unicolor*, and *C. a. yuracus*. *C. a. pleei* Hershkovitz 1949 and *C. a. leucocephalus* Gray 1865 (listed by Hershkovitz 1949) are regarded by Hernández-Camacho and Cooper (1976), Hernández Camacho and Defler (1991), and Defler (1994) as junior synonyms of *C. a. versicolor*. Bodini and Pérez-Hernández (1987) listed three subspecies for Venezuela: *C. a. adustus*, *C. a. leucocephalus* (regarded a junior synonym of *C. a. versicolor* by Hernández-Camacho and Cooper 1976), and *C. a. unicolor*. Bodini (1989) pointed out that Hill (1960) gives four subspecies for Venezuela, *C. a. cesarae*, *C. a. adustus*, *C. a. albifrons* and *C. a. unicolor*. Bodini and Pérez-Hernández (1987) give the form in the Federal Territory of Amazonas as *C. a. unicolor*, in agreement with Hershkovitz (1949), although Hershkovitz was uncertain of the limits of both this subspecies and *C. a. unicolor*. Hernández-Camacho and Cooper (1976) indicated that the Amazonian form in Venezuela was *C. a. albifrons* based on their belief that *C. a. unicolor* is a junior synonym. All subspecies of *C. albifrons* listed here are regarded as synonyms of just one species, *C. albifrons*, by Groves (1993).

9. Hill (1960) lists five subspecies of *Cebus capucinus*. *C. c. nigripectus* from the upper Río Cauca in Colombia, *C. c. capucinus* (Colombia), *C. c. imitator* (Costa Rica, including Coiba Island, and Panama), *C. c. limitaneus* (Honduras and Nicaragua), and *C. c. curtus* (Colombia). Hernández-Camacho and Cooper (1976) argued that variability in populations of these capuchins on the upper Río Cauca argued against the validity of *C. c. nigripectus*, and Hernández-Camacho and Defler (1991) and Defler (1994) listed just two subspecies of *C. capucinus* for Colombia: *C. c. capucinus* and *C. c. curtus*. Defler (1994) includes *C. c. nigripectus* under the name of *C. c. curtus*. *C. c. curtus* is endemic to the Colombian Island of Gorgona. It was possibly introduced in the 16th or 17th Centuries (Hernández-Camacho and Defler 1991) and is very similar to Panamanian *C. capucinus* (R. A. Mittermeier, pers. obs.). Silva-López *et al.* (1995) report on the possibility that *C. capucinus* may extend into Guatemala. Groves (1993) recognizes no subspecific differentiation in *C. capucinus*.

10. *Cebus olivaceus* requires a taxonomic revision. The subspecies listed are based on Hershkovitz (1949). The species name of *C. olivaceus* Schomburgk 1848 rather than *C. nigrivittatus* Wagner 1947 is argued by Husson (1978), and used by Eisenberg (1989). Groves (1993) gave all forms listed here as junior synonyms of *C. olivaceus*.

11. Bodini and Pérez-Hernández (1987) list five subspecies for Venezuela: *C. n. brunneus*, *C. n. apiculatus*, *C. n. nigrivittatus* (Amazonian), *C. n. olivaceus* and a fifth form

which has yet to be classified (widely distributed throughout central and northern Venezuela, to the north of Río Orinoco). Bodini (1989) pointed out that the subspecies of *C. nigritattus* are not well defined.

12. Taxonomy and distributions of *Pithecia* according to Hershkovitz (1987a) and Eisenberg (1989). Groves (1993) listed the following species: *P. aequatorialis*, *P. albicans*, *P. irrorata* (*P. i. vanzolinii* a synonym), *P. monachus* (*P. m. milleri* a synonym), *P. pithecia* (*P. p. chrysocephala* a synonym). A phylogeny at the generic level was recently proposed by Schneider *et al.* (1995).

13. *Chiropotes* taxonomy and distributions according to Hershkovitz (1985), Eisenberg (1989), and Ferrari (1995). Groves (1993) recognized just two species, *C. albinasus* and *C. satanas*: the subspecies of *C. satanas* recognized by Hershkovitz (1984) are considered by him to be synonyms.

14. Taxonomy and distributions of *Cacajao* according to Hernández-Camacho and Cooper (1976), Hershkovitz (1987b), Eisenberg (1989) and Boubli (1994). Groves (1989) listed the subspecies of *C. calvus* (*rubicundus*, *ucayalii*, and *novaesi*) recognized by Hershkovitz (1987b), as synonyms. Szalay and Delson (1979) gave *C. c. rubicundus* full species status.

15. Hill (1962) and Stanyon *et al.* (1995) list nine subspecific forms of *A. seniculus*: *A. s. seniculus*, *A. s. arctoidea*, *A. s. stramineus*, *A. s. macconelli*, *A. s. insulanus*, *A. s. amazonica*, *A. s. juara*, *A. s. puruensis*, *A. s. sara*. Cytogenetic studies have been carried out on; *A. s. seniculus* by Chu and Bender (1961), Bender and Chu (1963), and Yunis *et al.* (1976); *A. s. stramineus* by Lima and Seuánez (1991); *A. s. macconelli* by Lima *et al.* (1990); *A. s. sara* by Minezawa *et al.* (1985) and Stanyon *et al.* (1995); and *A. s. arctoidea* by Stanyon *et al.* (1995). *A. sara* is recognized here as a full species here following Minezawa *et al.* (1985; see also Groves 1993; Stanyon *et al.* 1995). Stanyon *et al.* (1995) concluded that the number of chromosomal differences between *A. s. sara* and *A. s. arctoidea* was on a similar scale to those found between *A. s. sara* and *A. s. seniculus* by Minezawa *et al.* (1985). *A. s. arctoidea* is here listed as a distinct species for this reason. Stanyon *et al.* (1995) noted that "as for other organisms in tropical forest regions, it is probable the biological diversity and number of species have been underestimated". Hill (1962) argued that the description of *A. s. juara* by Elliot (1910) confused the name and type locality. Hill (1962) demonstrated that it came in fact from the Río Juruá in Brazil, whereas Elliot gave the Río Juara in the Peruvian Amazon. Three years later, Elliot (1913) referred to the Río Juara in the Brazilian, not Peruvian, Amazon. Hill (1962) mentioned that *A. s. juara* is probably a junior synonym of *A. s. seniculus*.

16. An undescribed subspecies of *A. seniculus* north of the Río Orinoco, reported by Bodini and Pérez-Hernández (1987).

17. The taxonomy and distributions of *A. belzebul* follow Hill (1962), Langguth *et al.* (1987), and Bonvicino *et al.*

(1989). Cytogenetic studies have indicated, however, the probability that *A. b. nigerrima* is sufficiently distinct as to warrant species status (Armada *et al.* 1987; see also Lima and Seuánez 1989)).

18. The taxonomy and distribution of *Alouatta fusca* is discussed in Rylands *et al.* (1988; see also Hirsch *et al.* 1991). Occurrence of *A. fusca* in Argentina reviewed by Bitteti *et al.* (1994). Cytogenetic studies were carried out by Lima and Seuánez (1991).

19. Villalba *et al.* (1995) have raised the possibility that *A. caraya* extends as far south as Uruguay.

20. Taxonomic status and distribution of *Alouatta pigra* according to Smith (1970), Horwich (1983), Horwich and Johnson (1984), and Silva-López *et al.* (1995). *Alouatta palliata luctuosa* Lawrence, 1933 listed by Hill (1962) for Belize was not recognized by Froehlich and Froehlich (1986). *Alouatta pigra luctuosa* was listed by Dahl (1987) for the primates of Belize.

21. Groves (1993) lists the following species: *A. belzebul* (*discolor*, *nigerrima*, and *ululata* synonyms), *A. caraya*, *A. coibensis* (*trabeata* synonym), *A. fusca* (*clamitans* synonym), *A. palliata* (*aequatorialis*, *mexicana* synonyms), *A. pigra* (*A. p. luctuosa* synonym), *A. seniculus*, and *A. sara*. Distributions of *Alouatta* follow Hershkovitz (1949), Hill (1962), Froehlich and Froehlich (1986, 1987), Dahl (1987), Langguth *et al.* (1987), Bonvicino *et al.* (1989), Eisenberg (1989), Hirsch *et al.* (1991), Redford and Eisenberg (1992), and Curdts (1993). Note that Stanyon *et al.* (1995) gave the distribution of *A. seniculus amazonica* as the northern Atlantic forest, obviously an oversight rather than an affirmation. Their distribution map for the species is taken from Hill (1962) who placed *A. s. amazonica* in a small area north of the Río Solimões, to the west of the Río Negro, in the state of Amazonas, Brazil.

22. *Ateles* taxonomy according to Kellogg and Goldman (1944) and Konstant *et al.* (1985), except in the recognition of the forms *chamek* and *marginatus* as distinct species, following Groves (1989, 1993). De Boer and Bruijn (1990), Froehlich *et al.* (1991), and Medeiros (1994) argued that *A. paniscus* is a distinct form with no subspecies. De Boer and Bruijn (1990) indicated that *chamek* should be considered a full species. Froehlich *et al.* (1991) indicated that the form *chamek* is subspecific to *A. belzebuth* rather than *A. paniscus*. Hernández-Camacho and Cooper (1976), Hernández-Camacho and Defler (1991) refer to all *Ateles* as subspecies of *A. paniscus*, following Hershkovitz (1972). Defler (1994), on the other hand follows Kellogg and Goldman (1944) and refers to three species in Colombia: *A. belzebuth*, *A. fusciceps*, and *A. geoffroyi*. Groves (1989, 1993) gave the forms *chamek* Humboldt 1812, and *marginatus* É. Geoffroy 1809, species status, regarded *A. f. robustus* as a synonym of *A. f. fusciceps*, *A. b. brunneus* and *A. b. hybridus* as synonyms of *A. belzebuth*, and all *A. geoffroyi* subspecies as synonyms. Heltne and Kunkel (1975) provided additional taxonomic notes concerning pelage coloration in *A. paniscus* (including *chamek*) and *A.*

*fusciceps*. Kunkel *et al.* (1980), reported on chromosomal variation and zoogeography in the genus. Distributions according to Kellogg and Goldman (1944), Hershkovitz (1949), Hernández-Camacho and Cooper (1976), Martins *et al.* (1988), Eisenberg (1989), and Aquino and Encarnación (1994).

23. Hernández-Camacho and Defler (1985, 1991) and Defler (1994) also refer to *A. p. brunneus* Gray 1872 from the south-east of the Department of Bolívar, Colombia, listed here as a subspecies of *A. b. belzebuth* (see Hill 1962).

24. *Lagothrix* taxonomy and distributions according to Fooden (1963), Hernández-Camacho and Cooper (1976), Eisenberg (1989), Hernández-Camacho and Defler (1991) and Aquino and Encarnación (1994). This genus is in need of revision. Distribution of *L. flavicauda* follows Leo Luna (1987) and Butchart *et al.* (in press). Groves (1993) does not recognize the subspecific forms of *L. lagotricha*.

25. Vieira (1944) recognized two subspecies of *Brachyteles*. Recent evidence provided by Lemos de Sá *et al.* (1990, 1993), Fonseca *et al.* (1991) and Lemos de Sá and Glander (1993) indicates that Vieira's original (1944) standing was valid, but that differentiation is even more extreme and justifies the classification of the two forms as separate species. Distribution according to Aguirre (1971), Mittermeier *et al.* (1987), Santos *et al.* (1987), and Oliver and Santos (1991).

#### Mace-Lande Categories for Neotropical Primates

A draft version of the application of the Mace-Lande classification to the Neotropical primates was drawn up by the authors on the occasion of the XVth Congress of the International Primatological Society, held in Bali, 3-8 August 1994. This proposal was then sent to 86 members of the IUCN Species Survival Commission (SSC) Primate Specialist Group in October 1994 for comments and adjustments. The evaluation of the Mesoamerican primates also benefited from the concurrent preparation of the Primate Action Plan for the region (Rodríguez-Luna 1993). The final version is presented in Table 4, and Tables 5, 6, and 7 list the primates in each threatened category (Critically Endangered, Endangered and Vulnerable) with the criteria which determined their rating. Table 8 lists the species which are considered threatened according to the Mace-Lande System. Table 9 provides a summary of the number of species and taxa of Callitrichidae and Cebidae in the Critical (CR), Endangered (EN) and Vulnerable (VU), and the percentage of cebid and callitrichid and the total species and taxa in each threatened category is also shown.

From this evaluation it is possible to make the following statements:

#### Family Callitrichidae

1. This synthesis considers five genera, 35 species, and 55 species and subspecies of callitrichids.
2. Two of the five genera of callitrichids (40%) are threatened (*Leontopithecus* and *Callimico*).
3. Thirteen of the 35 species of callitrichids (37%) are threatened.
4. Sixteen of the 55 species and subspecies of callitrichids (29%) are threatened.
5. Seven of the 35 species of callitrichids (20%) are critically endangered or endangered (*Callithrix flaviceps*, *C. aurita*, *Saguinus oedipus*, *Leontopithecus rosalia*, *L. chrysomelas*, *L. chrysopygus*, *L. caissara*).
6. Eight of the 55 species and subspecies of callitrichids (14.5%) are critically endangered or endangered.

#### Family Cebidae

1. This synthesis considers 11 genera, 63 species, and 147 species and subspecies of cebids.
2. One of the 11 genera of cebids are threatened (*Brachyteles*).
3. Twenty of the 63 species of cebids (32%) are threatened.
4. Fifty-four of the 147 species and subspecies of the cebids (37%) are threatened.
5. Eight of the 63 species of cebids (13%) are critically endangered or endangered (*Aotus lemurinus*, *Saimiri oerstedii*, *Callicebus* sp., *Cebus xanthosternos*, *Alouatta coibensis*, *Ateles marginatus*, *Brachyteles arachnoides*, *B. hypoxanthus*).
6. Twenty-seven of the 147 species and subspecies of cebids (18%) are critically endangered or endangered.

#### New World Primates

1. This synthesis considers 16 genera, 98 species, and 202 species and subspecies of New World primates.
2. Three of the 16 genera of New World primates are threatened (*Callimico*, *Leontopithecus*, *Brachyteles*).
3. Thirty-three of the 98 species of New World primates (34%) are threatened.
4. Seventy of the 202 species and subspecies of

- New World primates (35%) are threatened.
5. Fifteen of 98 species of New World primates (15%) are critically endangered or endangered.
  6. Thirty-five of the 202 species and subspecies of New World primates (17%) are critically endangered or endangered.

#### **CRITICALLY ENDANGERED**

The two criteria applicable to all of the taxa listed as critically endangered taxa (Table 5) are: B. the extent of occurrence is estimated to be less than 100 km<sup>2</sup> or area of occupancy estimated to be less than 10 km<sup>2</sup>, and estimates indicating 1) Severely fragmented or known to exist at only a single location, and 2) Continuing decline, observed, inferred or projected in any of the following (a) extent of occurrence, (b) area of occupancy (c) area, extent and/or quality of habitat, (d) number of locations or subpopulations and (e) number of mature individuals. For all, except *Callicebus personatus barbarabrownae*, it was also possibly to invoke the following criterion: C. Population estimated to number less than 250 mature individuals and 2. A continuing decline, observed, projected, or inferred, in numbers of mature individuals

and population structure in the form of (a) severely fragmented (i.e., no subpopulation estimated to contain more than 50 mature individuals). For both *Alouatta fusca fusca* and *Leontopithecus caissara* it is also estimated that the population contains no more than 50 mature individuals (D).

#### **ENDANGERED**

Universal criteria for the endangered species and subspecies listed in Table 6 are the following: B. Extent of occurrence estimated to be less than 5000 km<sup>2</sup> or area of occupancy estimated to be less than 500 km<sup>2</sup>, and estimates indicating 1. Severely fragmented or known to exist at no more than five locations, and 2. Continuing decline, observed, inferred or projected, in any of the following: (a) extent of occurrence, (b) area of occupancy, (c) area, extent and/or quality of habitat, (d) number of locations or subpopulations and (e) number of mature individuals. In the case of all of the callitrichids, *Saimiri oerstedi oerstedi*, *Alouatta coibensis coibensis*, *Ateles geoffroyi griseescens*, *Ateles geoffroyi panamensis*, *Brachyteles arachnoides*, and *Brachyteles hypoxanthus* criterion C2 was also applied: C. Population estimated to number less than 2500 mature individuals and 2. A continuing decline,

Table 4. The IUCN Mace-Lande Categories for the status of New World Monkeys. Also shown is the distribution by country.

Species	Category	Distribution
<b>Family Callitrichidae</b>		
<i>Cebuella pygmaea</i>	LR	Bolivia, Brazil, Colombia, Ecuador, Peru
<i>Callithrix argentata</i>	LR	Brazil
<i>Callithrix leucippe</i>	VU	Brazil
<i>Callithrix melanura</i>	LR	Bolivia, Brazil, Paraguay
<i>Callithrix intermedia</i>	LR	Brazil
<i>Callithrix emiliae</i>	LR	Brazil
<i>Callithrix nigriceps</i>	VU	Brazil
<i>Callithrix marcai</i>	LR	Brazil
<i>Callithrix humeralifer</i>	LR	Brazil
<i>Callithrix chrysoleuca</i>	VU	Brazil
<i>Callithrix maeusi</i>	LR	Brazil
<i>Callithrix saterei</i>	DD	Brazil
<i>Callithrix jacchus</i>	LR	Brazil
<i>Callithrix penicillata</i>	LR	Brazil
<i>Callithrix kuhli</i>	LR	Brazil
<i>Callithrix geoffroyi</i>	VU	Brazil
<i>Callithrix flaviceps</i>	EN	Brazil
<i>Callithrix aurita</i>	EN	Brazil
<i>Saguinus nigricollis nigricollis</i>	LR	Brazil (?), Colombia, Peru
<i>Saguinus nigricollis graellsii</i>	LR	Colombia, Ecuador, Peru
<i>Saguinus nigricollis hernandezi</i>	VU	Colombia
<i>Saguinus fuscicollis fuscicollis</i>	LR	Brazil, Peru
<i>Saguinus fuscicollis fuscus</i>	LR	Brazil, Colombia

*continued ...*

Table 4 (continued)

<i>Saguinus fuscicollis avilapiresi</i>	LR	Brazil
<i>Saguinus fuscicollis cruzlimai</i>	DD	Brazil (?)
<i>Saguinus fuscicollis illigeri</i>	LR	Peru
<i>Saguinus fuscicollis leucogenys</i>	LR	Peru
<i>Saguinus fuscicollis nigrifrons</i>	LR	Peru
<i>Saguinus fuscicollis lagonotus</i>	LR	Ecuador, Peru
<i>Saguinus fuscicollis weddelli</i>	LR	Bolivia, Brazil, Peru
<i>Saguinus fuscicollis primitivus</i>	LR	Brazil
<i>Saguinus fuscicollis melanoleucus</i>	LR	Brazil
<i>Saguinus fuscicollis crandalli</i>	DD	Peru (?)
<i>Saguinus tripartitus</i>	LR	Ecuador, Peru
<i>Saguinus mystax mystax</i>	LR	Brazil, Peru
<i>Saguinus mystax pileatus</i>	LR	Brazil
<i>Saguinus mystax pluto</i>	LR	Brazil
<i>Saguinus labiatus labiatus</i>	LR	Bolivia, Brazil, Peru
<i>Saguinus labiatus thomasi</i>	LR	Brazil
<i>Saguinus imperator imperator</i>	VU	Brazil, Peru
<i>Saguinus imperator subgrisescens</i>	LR	Bolivia, Brazil, Peru
<i>Saguinus inustus</i>	LR	Brazil, Colombia
<i>Saguinus midas midas</i>	LR	Brazil, French Guiana, Guyana, Suriname
<i>Saguinus midas niger</i>	LR	Brazil
<i>Saguinus bicolor bicolor</i>	EN	Brazil
<i>Saguinus bicolor ochraceus</i>	LR	Brazil
<i>Saguinus bicolor martinsi</i>	LR	Brazil
<i>Saguinus leucopus</i>	VU	Colombia
<i>Saguinus oedipus</i>	EN	Colombia
<i>Saguinus geoffroyi</i>	LR	Colombia, Costa Rica, Panama
<i>Leontopithecus rosalia</i>	CR	Brazil
<i>Leontopithecus chrysomelas</i>	EN	Brazil
<i>Leontopithecus chrysopygus</i>	CR	Brazil
<i>Leontopithecus caissara</i>	CR	Brazil
<i>Callimico goeldii</i>	VU	Bolivia, Brazil, Colombia, Peru
<b>Family Cebidae</b>		
<i>Aotus lemurinus lemurinus</i>	VU	Colombia, Costa Rica, Panama
<i>Aotus lemurinus griseimembra</i>	EN	Colombia
<i>Aotus vociferans</i>	LR	Colombia, Brazil, Ecuador, Peru
<i>Aotus trivirgatus</i>	LR	Brazil, Colombia, Venezuela
<i>Aotus brumbacki</i>	VU	Colombia
<i>Aotus miconax</i>	VU	Peru
<i>Aotus nigriceps</i>	LR	Brazil, Colombia, Peru
<i>Aotus infuscatus</i>	LR	Brazil
<i>Aotus azarai azarai</i>	LR	Argentina, Bolivia, Paraguay
<i>Aotus azarai boliviensis</i>	LR	Bolivia, Peru
<i>Aotus nancymaae</i>	LR	Brazil, Colombia, Peru
<i>Aotus hershkovitzi</i>	DD	Colombia
<i>Callicebus modestus</i>	LR	Bolivia
<i>Callicebus donacophilus donacophilus</i>	LR	Bolivia
<i>Callicebus donacophilus pallescens</i>	LR	Bolivia, Brazil, Paraguay
<i>Callicebus olallae</i>	DD	Bolivia
<i>Callicebus oenanthe</i>	VU	Peru
<i>Callicebus cinerascens</i>	LR	Brazil
<i>Callicebus hoffmannsi hoffmannsi</i>	LR	Brazil
<i>Callicebus hoffmannsi baptista</i>	LR	Brazil
<i>Callicebus moloch</i>	LR	Brazil
<i>Callicebus brunneus</i>	LR	Brazil, Peru
<i>Callicebus cupreus cupreus</i>	LR	Brazil, Peru

continued ...

Table 4 (continued)

<i>Callicebus cupreus discolor</i>	LR	Colombia, Ecuador, Peru
<i>Callicebus cupreus ornatus</i>	VU	Colombia
<i>Callicebus caligatus</i>	LR	Brazil, Peru
<i>Callicebus dubius</i>	VU	Brazil
<i>Callicebus personatus personatus</i>	VU	Brazil
<i>Callicebus personatus nigrifrons</i>	VU	Brazil
<i>Callicebus personatus melanochir</i>	VU	Brazil
<i>Callicebus personatus barbarabrowniae</i>	CR	Brazil
<i>Callicebus</i> sp.	CR	Brazil
<i>Callicebus torquatus torquatus</i>	LR	Brazil
<i>Callicebus torquatus medemi</i>	VU	Colombia
<i>Callicebus torquatus lugens</i>	LR	Brazil, Colombia, Venezuela
<i>Callicebus torquatus lucifer</i>	LR	Brazil, Colombia, Peru
<i>Callicebus torquatus purinus</i>	LR	Brazil
<i>Callicebus torquatus regulus</i>	LR	Brazil
<i>Saimiri boliviensis boliviensis</i>	LR	Bolivia, Brazil, Peru
<i>Saimiri boliviensis peruviensis</i>	LR	Peru
<i>Saimiri boliviensis pluvialis</i>	LR	Brazil
<i>Saimiri boliviensis jaburuensis</i>	LR	Brazil
<i>Saimiri vanzolinii</i>	VU	Brazil
<i>Saimiri sciureus sciureus</i>	LR	Brazil, French Guiana, Guyana, Suriname
<i>Saimiri sciureus macrodon</i>	LR	Brazil, Colombia, Ecuador, Peru
<i>Saimiri sciureus cassiquiarensis</i>	LR	Brazil, Colombia, Venezuela
<i>Saimiri sciureus albigena</i>	LR	Colombia
<i>Saimiri oerstedi oerstedi</i>	EN	Costa Rica, Panama
<i>Saimiri oerstedi citrinellus</i>	CR	Costa Rica
<i>Saimiri ustus</i>	LR	Brazil
<i>Cebus apella apella</i>	LR	Colombia, Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Cebus apella macrocephalus</i>	LR	Brazil, Peru
<i>Cebus apella maranonis</i>	LR	Brazil, Colombia, Ecuador, Peru
<i>Cebus apella pallidus</i>	LR	Bolivia, Peru
<i>Cebus apella peruanus</i>	LR	Brazil, Peru
<i>Cebus apella libidinosus</i>	LR	Brazil
<i>Cebus apella nigritus</i>	LR	Argentina, Brazil
<i>Cebus apella robustus</i>	VU	Brazil
<i>Cebus apella margaritae</i>	CR	Venezuela
<i>Cebus apella paraguayanus</i>	LR	Argentina, Brazil, Paraguay
<i>Cebus xanthosternos</i>	CR	Brazil
<i>Cebus albifrons albifrons</i>	LR	Colombia
<i>Cebus albifrons unicolor</i>	LR	Colombia, Bolivia, Brazil, Peru, Venezuela
<i>Cebus albifrons adustus</i>	DD	Venezuela
<i>Cebus albifrons aequatorialis</i>	DD	Ecuador
<i>Cebus albifrons cesarae</i>	DD	Colombia
<i>Cebus albifrons cuscinus</i>	DD	Bolivia, Brazil, Peru
<i>Cebus albifrons hypoleucus</i>	DD	Colombia
<i>Cebus albifrons malitiosus</i>	DD	Colombia
<i>Cebus albifrons trinitatis</i>	CR	Trinidad
<i>Cebus albifrons versicolor</i>	DD	Colombia
<i>Cebus albifrons yuracus</i>	DD	Colombia, Ecuador, Peru
<i>Cebus albifrons leucocephalus</i>	DD	Venezuela
<i>Cebus capucinus capucinus</i>	LR	Colombia
<i>Cebus capucinus limitaneus</i>	LR	Belize, Honduras, Nicaragua
<i>Cebus capucinus imitator</i>	LR	Costa Rica, Panama
<i>Cebus capucinus curtus</i>	VU	Gorgona Island, Colombia
<i>Cebus olivaceus olivaceus</i>	LR	Brazil, Guyana, Venezuela,
<i>Cebus olivaceus apiculatus</i>	LR	Venezuela
<i>Cebus olivaceus brunneus</i>	LR	Venezuela

continued ...

Table 4 (continued)

<i>Cebus olivaceus castaneus</i>	LR	French Guiana, Suriname
<i>Cebus olivaceus</i> ssp.	DD	Venezuela
<i>Cebus kaapori</i>	VU	Brazil
<i>Pithecia pithecia pithecia</i>	LR	Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Pithecia pithecia chryscephala</i>	LR	Brazil
<i>Pithecia monachus monachus</i>	LR	Brazil, Colombia, Ecuador, Peru
<i>Pithecia monachus milleri</i>	VU	Colombia
<i>Pithecia irrorata irrorata</i>	LR	Bolivia, Brazil, Peru
<i>Pithecia irrorata vanzolinii</i>	LR	Brazil
<i>Pithecia albicans</i>	LR	Brazil
<i>Pithecia aequatorialis</i>	LR	Colombia (?), Ecuador, Peru
<i>Chiropotes albinasus</i>	LR	Brazil
<i>Chiropotes satanas satanas</i>	EN	Brazil
<i>Chiropotes satanas chiropotes</i>	LR	Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Chiropotes satanas utahicki</i>	VU	Brazil
<i>Cacajao calvus calvus</i>	EN	Brazil
<i>Cacajao calvus ucayalii</i>	VU	Brazil (?), Peru
<i>Cacajao calvus novaeisi</i>	EN	Brazil
<i>Cacajao calvus rubicundus</i>	EN	Brazil, Colombia
<i>Cacajao melanocephalus melanocephalus</i>	LR	Brazil, Venezuela
<i>Cacajao melanocephalus ouakary</i>	LR	Brazil, Colombia
<i>Alouatta seniculus seniculus</i>	LR	Colombia, Bolivia, Brazil, Ecuador, Peru, Venezuela
<i>Alouatta seniculus insulanus</i>	VU	Trinidad
<i>Alouatta seniculus stramineus</i>	LR	Brazil, Venezuela
<i>Alouatta seniculus macconnelli</i>	LR	Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Alouatta seniculus amazonica</i>	DD	Brazil
<i>Alouatta seniculus juara</i>	DD	Brazil
<i>Alouatta seniculus puruensis</i>	DD	Bolivia, Brazil
<i>Alouatta seniculus</i> ssp.	LR	Venezuela
<i>Alouatta arctoidea</i>	LR	Venezuela
<i>Alouatta sara</i>	LR	Bolivia
<i>Alouatta belzebul belzebul</i>	LR	Brazil
<i>Alouatta belzebul nigerrima</i>	LR	Brazil
<i>Alouatta belzebul discolor</i>	LR	Brazil
<i>Alouatta belzebul ululata</i>	CR	Brazil
<i>Alouatta fusca fusca</i>	CR	Brazil
<i>Alouatta fusca clamitans</i>	VU	Argentina, Brazil
<i>Alouatta palliata palliata</i>	LR	Costa Rica, Nicaragua, Honduras, Panama
<i>Alouatta palliata mexicana</i>	VU	Mexico, Guatemala
<i>Alouatta palliata aequatorialis</i>	LR	Colombia, Ecuador, Panama, Peru
<i>Alouatta coibensis coibensis</i>	EN	Coiba Island, Panama
<i>Alouatta coibensis trabeata</i>	CR	Panama
<i>Alouatta caraya</i>	LR	Argentina, Brazil, Paraguay, Uruguay (?)
<i>Alouatta pigra</i>	LR	Belize, Guatemala, Mexico
<i>Ateles geoffroyi geoffroyi</i>	LR	Nicaragua
<i>Ateles geoffroyi azuerensis</i>	CR	Panama
<i>Ateles geoffroyi frontatus</i>	VU	Costa Rica, Nicaragua
<i>Ateles geoffroyi griseescens</i>	EN	Colombia, Panama
<i>Ateles geoffroyi pan</i>	DD	Guatemala
<i>Ateles geoffroyi panamensis</i>	EN	Costa Rica, Panama

continued ...

Table 4 (continued)

<i>Ateles geoffroyi ornatus</i>	VU	Costa Rica
<i>Ateles geoffroyi vellerosus</i>	LR	El Salvador, Honduras, Guatemala, Mexico
<i>Ateles geoffroyi yucatanensis</i>	VU	Belize, Guatemala, Mexico
<i>Ateles fusciceps fusciceps</i>	CR	Ecuador
<i>Ateles fusciceps robustus</i> (= <i>rufiventris</i> )	VU	Colombia, Panama
<i>Ateles paniscus</i>	LR	Brazil, French Guiana, Guyana, Suriname
<i>Ateles belzebuth belzebuth</i>	VU	Brazil, Colombia, Ecuador, Peru, Venezuela
<i>Ateles belzebuth hybridus</i>	EN	Colombia, Venezuela
<i>Ateles belzebuth brunneus</i>	EN	Colombia
<i>Ateles chamek</i>	LR	Bolivia, Brazil, Peru
<i>Ateles marginatus</i>	EN	Brazil
<i>Lagothrix lagotricha lagotricha</i>	LR	Brazil, Colombia, Ecuador, Peru
<i>Lagothrix lagotricha cana</i>	VU	Brazil, Peru
<i>Lagothrix lagotricha poeppigii</i>	VU	Brazil, Ecuador, Peru
<i>Lagothrix lagotricha lugens</i>	CR	Colombia, Venezuela
<i>Lagothrix flavicauda</i>	CR	Peru
<i>Brachyteles arachnoides</i>	EN	Brazil
<i>Brachyteles hypoxanthus</i>	EN	Brazil

observed, projected, or inferred, in numbers of mature individuals and population structure in the form: (a) severely fragmented (i.e. no subpopulation estimated to contain more 250 mature individuals).

#### VULNERABLE

The criteria used for the categorization of the species and subspecies which are considered vulnerable (Table 7) are more varied, and include the following:

A. Population reduction in the form of the following : 1. An observed, estimated, inferred or suspected reduction of at least 20% over the last 10 years or three generations, whichever is the longer, based on (and specifying) any of the following : (c) a decline in area of occupancy, extent of occurrence and/or quality of habitat; B. Extent of occurrence estimated to be less than 20,000 km<sup>2</sup> or area of occupancy estimated to be less than 2000 km<sup>2</sup>, and estimates indicating any two of the following : 1. Severely fragmented or

Table 5. The New World primates (5 species and 16 taxa) considered to be Critically Endangered (CR) following the IUCN Mace-Lande Categories.

Species	Criteria	Distribution
<b>Family Callitrichidae</b>		
<i>Leontopithecus rosalia</i>	B1, B2, C2(a)	Brazil
<i>Leontopithecus chrysopygus</i>	B1, B2, C2(a)	Brazil
<i>Leontopithecus caissara</i>	B1, B2, C2(a), D	Brazil
<b>Family Cebidae</b>		
<i>Callicebus personatus barbarabrownae</i>	B1, B2	Brazil
<i>Callicebus</i> sp.	B1, B2, C2(a)	Brazil
<i>Saimiri oerstedi citrinellus</i>	B1, B2, C2(a)	Costa Rica
<i>Cebus apella margaritae</i>	B1, B2, C2(a)	Margarita Island, Venezuela
<i>Cebus albifrons trinitatis</i>	B1, B2, C2(a)	Trinidad
<i>Cebus xanthosternos</i>	B1, B2, C2(a)	Brazil
<i>Alouatta belzebul ululata</i>	B1, B2, C2(a)	Brazil
<i>Alouatta fusca fusca</i>	B1, B2, C2(a), D	Brazil
<i>Alouatta coibensis trabeata</i>	B1, B2, C2(a)	Panama
<i>Ateles geoffroyi azuerensis</i>	B1, B2, C2(a)	Panama
<i>Ateles fusciceps fusciceps</i>	B1, B2, C2(a)	Ecuador
<i>Lagothrix lagotricha lugens</i>	B1, B2, C2(a)	Colombia, Venezuela
<i>Lagothrix flavicauda</i>	B1, B2, C2(a)	Peru

known to exist at no more than ten locations. 2. Continuing decline, observed, inferred or projected, in either (a) extent of occurrence, (b) area of occupancy, (c) area, extent and/or quality of habitat, (d) number of locations or subpopulations, and (e) number of mature individuals; and C. Population estimated to number less than 10,000 mature individuals and 2. A continuing decline, observed, projected, or inferred, in numbers of mature individuals and populations structure in the form of either : (a)severely fragmented (i.e. no subpopulation estimated to contain more 1000 mature individuals) or (b) all individuals are in a single subpopulation.

in Table 32. From this evaluation it is possible to state the following:

#### Distribution by country

1. Neotropical primates are distributed through 20, possibly 21, countries from southern Mexico through Central America to Argentina and possibly Uruguay.

#### Family Callitrichidae

2. Callitrichids are distributed through 11 countries. That with the most species of callitrichids

Table 6. The New World primates considered to be Endangered (EN), following the IUCN Mace-Lande Categories.

Species	Criteria	Distribution
<b>Family Callitrichidae</b>		
<i>Callithrix flaviceps</i>	B1, B2, C2(a)	Brazil
<i>Callithrix aurita</i>	B1, B2, C2(a)	Brazil
<i>Saguinus bicolor bicolor</i>	B1, B2, C2(a)	Brazil
<i>Saguinus oedipus</i>	B1, B2, C2(a)	Colombia
<i>Leontopithecus chrysomelas</i>	B1, B2, C2(a)	Brazil
<b>Family Cebidae</b>		
<i>Aotus lemurinus griseimembra</i>	B1, B2	Colombia
<i>Saimiri oerstedi oerstedi</i>	B1, B2, C2(a)	Costa Rica, Panama
<i>Chiropotes satanas satanas</i>	B1, B2	Brazil
<i>Cacajao calvus calvus</i>	B1, B2	Brazil
<i>Cacajao calvus novaesi</i>	B1, B2	Brazil
<i>Cacajao calvus rubicundus</i>	B1, B2	Brazil, Colombia
<i>Alouatta coibensis coibensis</i>	B1, B2, C2(a)	Coiba Island, Panama
<i>Ateles geoffroyi griseescens</i>	B1, B2, C2 (a)	Colombia, Panama
<i>Ateles geoffroyi panamensis</i>	B1, B2, C2 (a)	Costa Rica, Panama
<i>Ateles marginatus</i>	B1, B2	Brazil
<i>Ateles belzebuth hybridus</i>	B1, B2	Colombia, Venezuela
<i>Ateles belzebuth brunneus</i>	B1, B2	Colombia
<i>Brachyteles arachnoides</i>	B1, B2, C2 (a)	Brazil
<i>Brachyteles hypoxanthus</i>	B1, B2, C2 (a)	Brazil

#### Primate Diversity by Country - Richness, Endemism, and Threatened Species

The primates occurring in each of the 21 countries, ranging from Mexico to northern Argentina and possibly Uruguay, are shown in Tables 10 to 30. Endemic species are in bold. A summary is provided in Table 31 and Figure 1 which give the number of species and taxa of callitrichids and cebids and the numbers for both families combined, along with the number of endemic species and taxa in each country. The numbers of threatened taxa in each country are shown

is Brazil (31 species and 44 species and subspecies). In second place is Peru, with eight species and 16 species and subspecies, and in third place is Colombia, with eight species and 10 species and subspecies. Six species occur in Bolivia, and the remaining seven countries have four or less species.

#### Family Cebidae

3. Cebids occur in 20, possibly 21 countries. That with the most species is Brazil, with

45 species and 84 species and subspecies. Colombia has 23 species and 41 species and subspecies. Third place is taken by Peru with 24 species and 35 species and subspecies. Venezuela is in fourth place with 13 species and 24 species and subspecies. Ecuador has 12 species and 14 species and subspecies and Bolivia has 11 species and 15 species and subspecies.

#### New World Primates

4. The country with the most primates is Brazil with 76 species (77% of the New World primate species) and 128 species and subspecies (63% of the New World primate taxa). Brazil is followed by Colombia with 31 species (32%) and 51 species and subspecies (25%) and Peru with 32 species (33%) and 51 species.

Table 7. The New World primates considered to be Vulnerable (VU), following the IUCN Mace-Lande Categories.

Species	Criteria	Distribution
<b>Family Callitrichidae</b>		
<i>Callithrix leucippe</i>	B2	Brazil
<i>Callithrix chrysoleuca</i>	B2	Brazil
<i>Callithrix nigriceps</i>	B2	Brazil
<i>Callithrix geoffroyi</i>	B1, B2(b), C2	Brazil
<i>Saguinus nigricollis hernandezi</i>	A1(c), B2	Colombia
<i>Saguinus imperator imperator</i>	A1(c), B2	Brazil, Peru
<i>Saguinus leucopus</i>	A1(c), B1, B2, C2	Colombia
<i>Callimico goeldii</i>	A1(c)	Bolivia, Brazil, Colombia, Peru
<b>Family Cebidae</b>		
<i>Aotus lemurinus lemurinus</i>	B1, B2, C2	Colombia, Costa Rica, Panama
<i>Aotus brumbacki</i>	B1, B2	Colombia
<i>Aotus miconax</i>	A1(c), B1, B2	Peru
<i>Callicebus oenanthe</i>	B1, B2	Peru
<i>Callicebus cupreus ornatus</i>	A1(c), B1, B2	Colombia
<i>Callicebus dubius</i>	B1, B2	Brazil
<i>Callicebus personatus personatus</i>	A1(c), B1, B2	Brazil
<i>Callicebus personatus nigrifrons</i>	A1(c)	Brazil
<i>Callicebus personatus melanochir</i>	A1(c)	Brazil
<i>Callicebus torquatus medemi</i>	B1, B2, C2	Colombia
<i>Saimiri vanzolinii</i>	B1, B2, C2	Brazil
<i>Cebus apella robustus</i>	B1, B2	Brazil
<i>Cebus kaapori</i>	A1(c), B1, B2	Brazil
<i>Cebus capucinus curtus</i>	B1, B2	Gorgona Island, Colombia
<i>Pithecia monachus milleri</i>	A1(c), B1, B2	Colombia
<i>Chiropotes satanas utahicki</i>	A1(c)	Brazil
<i>Cacajao calvus ucayalii</i>	A1(c)	Brazil (?), Peru
<i>Alouatta seniculus insulanus</i>	A1(c), B1, B2	Trinidad
<i>Alouatta fusca clamitans</i>	A1(c)	Argentina, Brazil
<i>Alouatta palliata mexicana</i>	A1(c), B1, B2	Mexico, Guatemala
<i>Ateles geoffroyi frontatus</i>	A1(c), B1, B2	Costa Rica, Nicaragua
<i>Ateles geoffroyi ornatus</i>	A1(c), B1, B2	Costa Rica
<i>Ateles geoffroyi yucatanensis</i>	A1(c), B1, B2	Belize, Guatemala, Mexico
<i>Ateles fusciceps robustus (= rufiventris)</i>	A1(c), B1, B2	Colombia, Panama
<i>Ateles belzebuth belzebuth</i>	A1(c)	Brazil, Colombia, Ecuador, Peru, Venezuela
<i>Lagothrix lagotricha cana</i>	A1(c)	Brazil, Peru
<i>Lagothrix lagotricha poeppigii</i>	A1(c)	Brazil, Ecuador, Peru

Table 8. Threatened species of Neotropical Primates.

Species
<b>Family Callitrichidae</b>
<i>Callithrix leucippe</i>
<i>Callithrix chrysoleuca</i>
<i>Callithrix nigriceps</i>
<i>Callithrix geoffroyi</i>
<i>Callithrix flaviceps</i>
<i>Callithrix aurita</i>
<i>Saguinus leucopus</i>
<i>Saguinus oedipus</i>
<i>Leontopithecus rosalia</i>
<i>Leontopithecus chrysomelas</i>
<i>Leontopithecus chrysopygus</i>
<i>Leontopithecus caissara</i>
<i>Callimico goeldii</i>
<b>Family Cebidae</b>
<i>Aotus lemurinus</i>
<i>Aotus brumbacki</i>
<i>Aotus miconax</i>
<i>Callicebus oenanthae</i>
<i>Callicebus dubius</i>
<i>Callicebus personatus</i>
<i>Callicebus sp.</i>
<i>Saimiri vanzolinii</i>
<i>Saimiri oerstedi</i>
<i>Cebus xanthosternos</i>
<i>Cebus kaapori</i>
<i>Cacajao calvus</i>
<i>Alouatta fusca</i>
<i>Alouatta coibensis</i>
<i>Ateles fusciceps</i>
<i>Ateles belzebuth</i>
<i>Ateles marginatus</i>
<i>Lagothrix flavicauda</i>
<i>Brachyteles arachnoides</i>
<i>Brachyteles hypoxanthus</i>




Figure 1. Number of Neotropical primate species and genera by country, with number of endemics indicated (map by R. B. Machado).

cies and subspecies (25%). Bolivia has 17 species and 21 species and subspecies, Venezuela has 13 species and 24 species and subspecies, and Ecuador 16 species and 18 species and subspecies. Panama has the most primates in Central America, with eight species and 12 species and subspecies.

#### Endemism

5. Fifty species, or 51% of the Neotropical primate species, are endemic to one country. A total of 114 species and subspecies, or 56% of the Neotropical primate taxa, are restricted to just one country. Six countries have endemic species: Brazil (38), Colombia (6), Bolivia (3),

Table 9. A summary of the number of species and taxa of Callitrichidae and Cebidae in the IUCN Mace-Lande categories of Critical (CR), Endangered (EN) and Vulnerable (VU). The percentage of cebid and callitrichid and the total species and taxa in each threatened category is also shown.

Family	Genus	Species	Threatened Category (sp.)				Threatened Category (taxa)				
			CR	EN	VU	Total	CR	EN	VU	Total	
Callitrichidae	5	35	55	3	4	6	13	3	5	8	16
				8.6%	11.4%	17.1%	37.1%	5.4%	9.1%	14.5%	29.1%
Cebidae	11	63	147	3	3	6	20	13	14	27	54
				4.8%	4.8%	9.7%	31.7%	8.8%	9.5%	18.4%	36.7%
<b>TOTAL</b>		<b>16</b>	<b>98</b>	<b>202</b>				<b>16</b>	<b>19</b>	<b>35</b>	<b>69</b>
				6.1%	7.2%	12.4%	33.7%	7.9%	9.4%	17.3%	34.1%

Peru (3), Ecuador (2), and Panama (1). Venezuela has six endemic subspecies and one endemic species. Costa Rica and Trinidad each have two endemic subspecies.

#### *Threatened Species by Country*

6. Brazil has the highest number of threatened species and subspecies 38 taxa accounting for 29% of its primates. Thirteen of 44 callitrichids (30%) are threatened, 25 of Brazil's 84 cebids (30%) are threatened. Brazil has 69 endemic taxa, 30 (43%) of which are threatened.
7. Colombia has 18 threatened species and subspecies which account for 35% of its 51 primates. Colombia has 18 endemic taxa, 10 (55%) of which are threatened.
8. Peru has ten threatened species and subspecies which account for 20% of its 51 primates. Peru has seven endemic taxa, three (43%) of which are threatened.
9. Panama has eight threatened species and subspecies which account for two-thirds of its primates. All three of Panama's endemic primates are threatened.
10. Costa Rica has six threatened species and subspecies, accounting for two-thirds of its primates. The two primates endemic to Costa Rica are threatened.
11. Venezuela has four threatened species and subspecies, accounting for 17% of its primates. None of its seven endemic taxa are ranked as threatened.
12. Ecuador has three threatened species and subspecies, accounting for 18% of its primates. One of Ecuador's two endemic primates is threatened.
13. Mexico has two threatened taxa, accounting for one-half of its primates. Guatemala has two threatened taxa.
14. Argentina, Belize, Bolivia, Honduras, Nicaragua and Trinidad each have one threatened taxon.
15. Trinidad has two threatened endemic primates
16. No threatened species or subspecies are recorded for El Salvador, French Guiana, Guyana, Paraguay, Suriname, and Uruguay.

#### **ARGENTINA**

Table 10. The IUCN Mace-Lande Categories for the conservation status of the primates of Argentina.

Species	Category	Distribution
<b>Family Cebidae</b>		
<i>Aotus azarae azarae</i>	LR	Argentina, Bolivia, Paraguay
<i>Cebus apella nigritus</i>	LR	Argentina, Brazil
<i>Cebus apella paraguayanus</i>	LR	Argentina, Brazil, Paraguay
<i>Alouatta fusca clamitans</i>	VU	Argentina, Brazil
<i>Alouatta caraya</i>	LR	Argentina, Brazil, Paraguay, Uruguay (?)

#### **BELIZE**

Table 11. The IUCN Mace-Lande Categories for the conservation status of the primates of Belize.

Species	Category	Distribution
<b>Family Cebidae</b>		
<i>Cebus capucinus limitaneus</i>	LR	Belize, Honduras, Nicaragua
<i>Alouatta pigra</i>	LR	Belize, Guatemala, Mexico
<i>Ateles geoffroyi yucatanensis</i>	VU	Belize, Guatemala, Mexico

**BOLIVIA**

Table 12. The IUCN Mace-Lande Categories for the conservation status of the primates of Bolivia. Endemic taxa are in bold.

Species	Category	Distribution
<b>Family Callitrichidae</b>		
<i>Cebuella pygmaea</i>	LR	Bolivia, Brazil, Colombia, Ecuador, Peru
<i>Callithrix melanura</i>	LR	Bolivia, Brazil, Paraguay
<i>Saguinus fuscicollis weddelli</i>	LR	Bolivia, Brazil, Peru
<i>Saguinus labiatus labiatus</i>	LR	Bolivia, Brazil, Peru
<i>Saguinus imperator subgriseus</i>	LR	Bolivia, Brazil, Peru
<i>Callimico goeldii</i>	VU	Bolivia, Brazil, Colombia, Peru
<b>Family Cebidae</b>		
<i>Aotus azarae azarae</i>	LR	Argentina, Bolivia, Paraguay
<i>Aotus azarae boliviensis</i>	LR	Bolivia, Peru
<i>Callicebus modestus</i>	LR	<b>Bolivia</b>
<i>Callicebus donacophilus donacophilus</i>	LR	<b>Bolivia</b>
<i>Callicebus donacophilus pallescens</i>	LR	Bolivia, Brazil, Paraguay
<i>Callicebus olallae</i>	DD	<b>Bolivia</b>
<i>Saimiri boliviensis boliviensis</i>	LR	Bolivia, Brazil, Peru
<i>Cebus apella pallidus</i>	LR	Bolivia, Peru
<i>Cebus albifrons unicolor</i>	LR	Colombia, Bolivia, Brazil, Peru, Venezuela
<i>Cebus albifrons cuscinus</i>	DD	Bolivia, Brazil, Peru
<i>Pithecia irrorata irrorata</i>	LR	Bolivia, Brazil, Peru
<i>Alouatta seniculus seniculus</i>	LR	Colombia, Bolivia, Brazil, Ecuador, Peru, Venezuela
<i>Alouatta seniculus puruensis</i>	LR	Bolivia, Brazil
<i>Alouatta sara</i>	LR	<b>Bolivia</b>
<i>Ateles chamek</i>	LR	Bolivia, Brazil, Peru

**BRAZIL**

Table 13. The IUCN Mace-Lande Categories for the conservation status of the primates of Brazil. Endemic taxa are in bold.

Species	Category	Distribution
<b>Family Callitrichidae</b>		
<i>Cebuella pygmaea</i>	LR	Bolivia, Brazil, Colombia, Ecuador, Peru
<i>Callithrix argentata</i>	LR	<b>Brazil</b>
<i>Callithrix leucippe</i>	VU	<b>Brazil</b>
<i>Callithrix melanura</i>	LR	Bolivia, Brazil, Paraguay
<i>Callithrix intermedia</i>	LR	<b>Brazil</b>
<i>Callithrix emiliae</i>	LR	<b>Brazil</b>
<i>Callithrix nigriceps</i>	VU	<b>Brazil</b>
<i>Callithrix mariae</i>	LR	<b>Brazil</b>
<i>Callithrix humeralifer</i>	LR	<b>Brazil</b>
<i>Callithrix chrysoleuca</i>	VU	<b>Brazil</b>
<i>Callithrix maeusi</i>	LR	<b>Brazil</b>
<i>Callithrix saterei</i>	DD	<b>Brazil</b>
<i>Callithrix jacchus</i>	LR	<b>Brazil</b>
<i>Callithrix penicillata</i>	LR	<b>Brazil</b>

continued ...

Table 13 (continued)

<i>Callithrix kuhli</i>	LR	Brazil
<i>Callithrix geoffroyi</i>	VU	Brazil
<i>Callithrix flaviceps</i>	EN	Brazil
<i>Callithrix aurita</i>	EN	Brazil
<i>Saguinus nigricollis nigricollis</i>	LR	Brazil (?), Colombia, Peru
<i>Saguinus fuscicollis fuscicollis</i>	LR	Brazil, Peru
<i>Saguinus fuscicollis fuscus</i>	LR	Brazil, Colombia
<i>Saguinus fuscicollis avilapiresi</i>	LR	Brazil
<i>Saguinus fuscicollis cruzlimai</i>	DD	Brazil (?)
<i>Saguinus fuscicollis weddelli</i>	LR	Bolivia, Brazil, Peru
<i>Saguinus fuscicollis primitivus</i>	LR	Brazil
<i>Saguinus fuscicollis melanoleucus</i>	LR	Brazil
<i>Saguinus mystax mystax</i>	LR	Brazil, Peru
<i>Saguinus mystax pileatus</i>	LR	Brazil
<i>Saguinus mystax pluto</i>	LR	Brazil
<i>Saguinus labiatus labiatus</i>	LR	Bolivia, Brazil, Peru
<i>Saguinus labiatus thomasi</i>	LR	Brazil
<i>Saguinus imperator imperator</i>	VU	Brazil, Peru
<i>Saguinus imperator subgriseus</i>	LR	Bolivia, Brazil, Peru
<i>Saguinus inustus</i>	LR	Brazil, Colombia
<i>Saguinus midas midas</i>	LR	Brazil, French Guiana, Guyana, Suriname
<i>Saguinus midas niger</i>	LR	Brazil
<i>Saguinus bicolor bicolor</i>	EN	Brazil
<i>Saguinus bicolor ochraceus</i>	LR	Brazil
<i>Saguinus bicolor martinsi</i>	LR	Brazil
<i>Leontopithecus rosalia</i>	CR	Brazil
<i>Leontopithecus chrysomelas</i>	EN	Brazil
<i>Leontopithecus chrysopygus</i>	CR	Brazil
<i>Leontopithecus caissara</i>	CR	Brazil
<i>Callimico goeldii</i>	VU	Bolivia, Brazil, Colombia, Peru
<b>Family Cebidae</b>		
<i>Aotus vociferans</i>	LR	Colombia, Brazil, Ecuador, Peru
<i>Aotus trivirgatus</i>	LR	Brazil, Colombia, Venezuela
<i>Aotus nigriceps</i>	LR	Brazil, Colombia, Peru
<i>Aotus infuscatus</i>	LR	Brazil
<i>Aotus azarae azarae</i>	LR	Bolivia, Brazil, Paraguay
<i>Aotus nancymaae</i>	LR	Brazil, Colombia, Peru
<i>Callicebus donacophilus pallescens</i>	LR	Bolivia, Brazil, Paraguay
<i>Callicebus cinerascens</i>	LR	Brazil
<i>Callicebus hoffmannsi hoffmannsi</i>	LR	Brazil
<i>Callicebus hoffmannsi baptista</i>	LR	Brazil
<i>Callicebus moloch</i>	LR	Brazil
<i>Callicebus brunneus</i>	LR	Brazil, Peru
<i>Callicebus cupreus cupreus</i>	LR	Brazil, Peru
<i>Callicebus caligatus</i>	LR	Brazil, Peru
<i>Callicebus dubius</i>	VU	Brazil
<i>Callicebus personatus personatus</i>	VU	Brazil
<i>Callicebus personatus nigrifrons</i>	VU	Brazil
<i>Callicebus personatus melanochir</i>	VU	Brazil
<i>Callicebus personatus barbarabrownae</i>	CR	Brazil
<i>Callicebus sp.</i>	CR	Brazil
<i>Callicebus torquatus torquatus</i>	LR	Brazil
<i>Callicebus torquatus lugens</i>	LR	Brazil, Colombia, Venezuela
<i>Callicebus torquatus lucifer</i>	LR	Brazil, Colombia, Peru
<i>Callicebus torquatus purinus</i>	LR	Brazil
<i>Callicebus torquatus regulus</i>	LR	Brazil

continued ...

Table 13 (continued)

<i>Saimiri boliviensis boliviensis</i>	LR	Bolivia, Brazil, Peru
<i>Saimiri boliviensis pluvialis</i>	LR	Brazil
<i>Saimiri boliviensis jaburuensis</i>	LR	Brazil
<i>Saimiri vanzolinii</i>	VU	Brazil
<i>Saimiri sciureus sciureus</i>	LR	Brazil, French Guiana, Guyana, Suriname
<i>Saimiri sciureus macrodon</i>	LR	Brazil, Colombia, Ecuador, Peru
<i>Saimiri sciureus cassiquiarensis</i>	LR	Brazil, Colombia, Venezuela
<i>Saimiri ustus</i>	LR	Brazil
<i>Cebus apella apella</i>	LR	Colombia, Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Cebus apella macrocephalus</i>	LR	Brazil, Peru
<i>Cebus apella maranonis</i>	LR	Brazil, Colombia, Ecuador, Peru
<i>Cebus apella peruanus</i>	LR	Brazil, Peru
<i>Cebus apella libidinosus</i>	LR	Brazil
<i>Cebus apella nigritus</i>	LR	Argentina, Brazil
<i>Cebus apella robustus</i>	VU	Brazil
<i>Cebus apella paraguayanus</i>	LR	Argentina, Brazil, Paraguay
<i>Cebus xanthosternos</i>	CR	Brazil
<i>Cebus albifrons unicolor</i>	LR	Colombia, Bolivia, Brazil, Peru, Venezuela
<i>Cebus albifrons cuscinus</i>	DD	Bolivia, Brazil, Peru
<i>Cebus olivaceus olivaceus</i>	LR	Brazil, Guyana, Venezuela
<i>Cebus kaapori</i>	VU	Brazil
<i>Pithecia pithecia pithecia</i>	LR	Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Pithecia pithecia chrysocephala</i>	LR	Brazil
<i>Pithecia monachus monachus</i>	LR	Brazil, Colombia, Ecuador, Peru
<i>Pithecia irrorata irrorata</i>	LR	Bolivia, Brazil, Peru
<i>Pithecia irrorata vanzolinii</i>	LR	Brazil
<i>Pithecia albicans</i>	LR	Brazil
<i>Chiropotes albinasus</i>	LR	Brazil
<i>Chiropotes satanas satanas</i>	EN	Brazil
<i>Chiropotes satanas chiropotes</i>	LR	Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Chiropotes satanas utahicki</i>	VU	Brazil
<i>Cacajao calvus calvus</i>	EN	Brazil
<i>Cacajao calvus ucayalii</i>	VU	Brazil (?), Peru
<i>Cacajao calvus novaesi</i>	EN	Brazil
<i>Cacajao calvus rubicundus</i>	EN	Brazil, Colombia
<i>Cacajao melanocephalus melanocephalus</i>	LR	Brazil, Venezuela
<i>Cacajao melanocephalus ouakary</i>	LR	Brazil, Colombia
<i>Alouatta seniculus seniculus</i>	LR	Colombia, Bolivia, Brazil, Ecuador, Peru, Venezuela
<i>Alouatta seniculus stramineus</i>	LR	Brazil, Venezuela
<i>Alouatta seniculus macconnelli</i>	LR	Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Alouatta seniculus amazonica</i>	LR	Brazil
<i>Alouatta seniculus juara</i>	LR	Brazil, Peru
<i>Alouatta seniculus puruensis</i>	LR	Bolivia, Brazil
<i>Alouatta belzebul belzebul</i>	LR	Brazil
<i>Alouatta belzebul nigerrima</i>	LR	Brazil
<i>Alouatta belzebul discolor</i>	LR	Brazil
<i>Alouatta belzebul ululata</i>	CR	Brazil
<i>Alouatta fusca fusca</i>	CR	Brazil
<i>Alouatta fusca clamitans</i>	VU	Argentina, Brazil
<i>Alouatta caraya</i>	LR	Argentina, Brazil, Paraguay, Uruguay (?)
<i>Ateles paniscus</i>	LR	Brazil, French Guiana, Guyana, Suriname
<i>Ateles belzebuth belzebuth</i>	VU	Brazil, Colombia, Ecuador, Peru, Venezuela
<i>Ateles chamek</i>	LR	Bolivia, Brazil, Peru

continued ...

Table 13 (continued)

<b><i>Ateles marginatus</i></b>	<b>EN</b>	<b>Brazil</b>
<i>Lagothrix lagotricha lagotricha</i>	LR	Brazil, Colombia, Ecuador, Peru
<i>Lagothrix lagotricha cana</i>	VU	Brazil, Peru
<i>Lagothrix lagotricha poeppigii</i>	VU	Brazil, Ecuador, Peru
<b><i>Brachyteles arachnoides</i></b>	<b>EN</b>	<b>Brazil</b>
<b><i>Brachyteles hypoxanthus</i></b>	<b>EN</b>	<b>Brazil</b>

**COLOMBIA**

Table 14. The IUCN Mace-Lande Categories for the conservation status of the primates of Colombia. Endemic taxa are in bold.

Species	Category	Distribution
<b>Family Callitrichidae</b>		
<i>Cebuella pygmaea</i>	LR	Bolivia, Brazil, Colombia, Ecuador, Peru
<i>Saguinus nigricollis nigricollis</i>	LR	Brazil (?), Colombia, Peru
<i>Saguinus nigricollis graellsi</i>	LR	Colombia, Ecuador, Peru
<b><i>Saguinus nigricollis hernandezii</i></b>	VU	<b>Colombia</b>
<i>Saguinus fuscicollis fuscus</i>	LR	Brazil, Colombia
<i>Saguinus inustus</i>	LR	Brazil, Colombia
<b><i>Saguinus leucopus</i></b>	VU	<b>Colombia</b>
<b><i>Saguinus oedipus</i></b>	EN	<b>Colombia</b>
<i>Saguinus geoffroyi</i>	LR	Colombia, Costa Rica, Panama
<i>Callimico goeldii</i>	VU	Bolivia, Brazil, Colombia, Peru
<b>Family Cebidae</b>		
<i>Aotus lemurinus lemurinus</i>	VU	Colombia, Costa Rica, Panama
<b><i>Aotus lemurinus griseimembra</i></b>	EN	<b>Colombia</b>
<i>Aotus vociferans</i>	LR	Colombia, Brazil, Ecuador, Peru
<i>Aotus trivirgatus</i>	LR	Brazil, Colombia, Venezuela
<b><i>Aotus brumbacki</i></b>	VU	<b>Colombia</b>
<i>Aotus nigriceps</i>	LR	Brazil, Colombia, Peru
<i>Aotus nancymaae</i>	LR	Brazil, Colombia, Peru
<b><i>Aotus hershkovitzii</i></b>	DD	<b>Colombia</b>
<i>Callicebus cupreus discolor</i>	LR	Colombia, Ecuador, Peru
<b><i>Callicebus cupreus ornatus</i></b>	VU	<b>Colombia</b>
<b><i>Callicebus torquatus medemi</i></b>	VU	<b>Colombia</b>
<i>Callicebus torquatus lugens</i>	LR	Brazil, Colombia, Venezuela
<i>Callicebus torquatus lucifer</i>	LR	Brazil, Colombia, Peru
<i>Saimiri sciureus macrodon</i>	LR	Brazil, Colombia, Ecuador, Peru
<i>Saimiri sciureus cassiquiarensis</i>	LR	Brazil, Colombia, Venezuela
<b><i>Saimiri sciureus albigena</i></b>	LR	<b>Colombia</b>
<i>Cebus apella apella</i>	LR	Colombia, Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Cebus apella maranensis</i>	LR	Brazil, Colombia, Ecuador, Peru
<b><i>Cebus albifrons albifrons</i></b>	LR	<b>Colombia</b>
<i>Cebus albifrons unicolor</i>	LR	Colombia, Bolivia, Brazil, Peru, Venezuela
<i>Cebus albifrons cesareae</i>	DD	<b>Colombia</b>
<i>Cebus albifrons hypoleucus</i>	DD	<b>Colombia</b>
<i>Cebus albifrons malitiosus</i>	DD	<b>Colombia</b>
<i>Cebus albifrons versicolor</i>	DD	<b>Colombia</b>
<i>Cebus albifrons yuracus</i>	DD	Colombia, Ecuador, Peru

continued ...

Table 14 (continued)

<i>Cebus capucinus capucinus</i>	LR	<b>Colombia</b>
<i>Cebus capucinus curtus</i>	VU	Gorgona Island, Colombia
<i>Pithecia monachus monachus</i>	LR	Brazil, Colombia, Ecuador, Peru
<i>Pithecia monachus milleri</i>	VU	<b>Colombia</b>
<i>Pithecia aequatorialis</i>	LR	Colombia (?), Ecuador, Peru
<i>Cacajao calvus rubicundus</i>	EN	Brazil, Colombia
<i>Cacajao melanocephalus ouakary</i>	LR	Brazil, Colombia
<i>Alouatta seniculus seniculus</i>	LR	Colombia, Bolivia, Brazil, Ecuador, Peru, Venezuela
<i>Alouatta palliata aequatorialis</i>	LR	Colombia, Ecuador, Panama, Peru
<i>Ateles geoffroyi griseescens</i>	EN	Colombia, Panama
<i>Ateles fusciceps robustus (= rufiventris)</i>	VU	Colombia, Panama
<i>Ateles belzebuth belzebuth</i>	VU	Brazil, Colombia, Ecuador, Peru, Venezuela
<i>Ateles belzebuth hybridus</i>	EN	Colombia, Venezuela
<i>Ateles belzebuth brunneus</i>	EN	<b>Colombia</b>
<i>Lagothrix lagotricha lagotricha</i>	LR	Brazil, Colombia, Ecuador, Peru
<i>Lagothrix lagotricha lugens</i>	CR	Colombia, Venezuela

**COSTA RICA**

Table 15. The IUCN Mace-Lande Categories for the conservation status of the primates of Costa Rica. Endemic taxa are in bold.

Species	Category	Distribution
<b>Family Callitrichidae</b>		
<i>Saguinus geoffroyi</i>	LR	Colombia, Costa Rica, Panama
<b>Family Cebidae</b>		
<i>Aotus lemurinus lemurinus</i>	VU	Colombia, Costa Rica, Panama
<i>Saimiri oerstedi oerstedi</i>	EN	Costa Rica, Panama
<i>Saimiri oerstedi citrinellus</i>	CR	<b>Costa Rica</b>
<i>Cebus capucinus imitator</i>	LR	Costa Rica, Panama
<i>Alouatta palliata palliata</i>	LR	Costa Rica, Nicaragua, Honduras, Panama
<i>Ateles geoffroyi frontatus</i>	VU	Costa Rica, Nicaragua
<i>Ateles geoffroyi panamensis</i>	EN	Costa Rica, Panama
<i>Ateles geoffroyi ornatus</i>	VU	<b>Costa Rica</b>

**ECUADOR**

Table 16. The IUCN Mace-Lande Categories for the conservation status of the primates of Ecuador. Endemic taxa are in bold.

Species	Category	Distribution
<b>Family Callitrichidae</b>		
<i>Cebuella pygmaea</i>	LR	Bolivia, Brazil, Colombia, Ecuador, Perú
<i>Saguinus nigricollis graellsii</i>	LR	Colombia, Ecuador, Peru
<i>Saguinus fuscicollis lagonotus</i>	LR	Ecuador, Peru
<i>Saguinus tripartitus</i>	LR	Ecuador, Peru

continued ...

Table 16 (continued)

<b>Family Cebidae</b>		
<i>Aotus vociferans</i>	LR	Colombia, Brazil, Ecuador, Peru
<i>Callicebus cupreus discolor</i>	LR	Colombia, Ecuador, Peru
<i>Saimiri sciureus macrodon</i>	LR	Brazil, Colombia, Ecuador, Peru
<i>Cebus apella maranonis</i>	LR	Brazil, Colombia, Ecuador, Peru
<i>Cebus albifrons aequatorialis</i>	DD	Ecuador
<i>Cebus albifrons yuracu</i>	DD	Colombia, Ecuador, Peru
<i>Pithecia monachus monachus</i>	LR	Brazil, Colombia, Ecuador, Peru
<i>Pithecia aequatorialis</i>	LR	Colombia (?), Ecuador, Peru
<i>Alouatta seniculus seniculus</i>	LR	Colombia, Bolivia, Brazil, Ecuador, Peru, Venezuela
<i>Alouatta palliata aequatorialis</i>	LR	Colombia, Ecuador, Panama, Peru
<i>Atelos fusciceps fusciceps</i>	CR	Ecuador
<i>Atelos belzebuth belzebuth</i>	VU	Brazil, Colombia, Ecuador, Peru, Venezuela
<i>Lagothrix lagotricha lagotricha</i>	LR	Brazil, Colombia, Ecuador, Peru
<i>Lagothrix lagotricha poeppigii</i>	VU	Brazil, Ecuador, Peru

**EL SALVADOR**

Table 17. The IUCN Mace-Lande Categories for the conservation status of the primates of El Salvador.

Species	Category	Distribution
<b>Family Cebidae</b>		
<i>Atelos geoffroyi vellerosus</i>	LR	El Salvador, Honduras, Guatemala, Mexico

**FRENCH GUIANA**

Table 18. The IUCN Mace-Lande Categories for the conservation status of the primates of French Guiana.

Species	Category	Distribution
<b>Family Callitrichidae</b>		
<i>Saguinus midas midas</i>	LR	Brazil, French Guiana, Guyana, Suriname
<b>Family Cebidae</b>		
<i>Saimiri sciureus sciureus</i>	LR	Brazil, French Guiana, Guyana, Suriname
<i>Cebus apella apella</i>	LR	Colombia, Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Cebus olivaceus castaneus</i>	LR	French Guiana, Suriname
<i>Pithecia pithecia pithecia</i>	LR	Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Chiropotes satanas chiropotes</i>	LR	Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Alouatta seniculus macconnelli</i>	LR	Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Atelos paniscus</i>	LR	Brazil, French Guiana, Guyana, Suriname

**GUATEMALA**

Table 19. The IUCN Mace-Lande Categories for the conservation status of the primates of Guatemala.

Species	Category	Distribution
<b>Family Cebidae</b>		
<i>Alouatta palliata mexicana</i>	VU	Guatemala, Mexico
<i>Alouatta pigra</i>	LR	Belize, Guatemala, Mexico
<i>Ateles geoffroyi vellerosus</i>	LR	El Salvador, Honduras, Guatemala, Mexico
<i>Ateles geoffroyi yucatanensis</i>	VU	Belize, Guatemala, Mexico

**GUYANA**

Table 20. The IUCN Mace-Lande Categories for the conservation status of the primates of Guyana.

Species	Category	Distribution
<b>Family Callitrichidae</b>		
<i>Saguinus midas midas</i>	LR	Brazil, French Guiana, Guyana, Suriname
<b>Family Cebidae</b>		
<i>Saimiri sciureus sciureus</i>	LR	Brazil, French Guiana, Guyana, Suriname
<i>Cebus apella apella</i>	LR	Colombia, Brazil, French Guiana, Guyana, Suriname Venezuela
<i>Cebus olivaceus olivaceus</i>	LR	Brazil, Guyana, Venezuela,
<i>Pithecia pithecia pithecia</i>	LR	Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Chiropotes satanas chiropotes</i>	LR	Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Alouatta seniculus macconnelli</i>	LR	Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Ateles paniscus</i>	LR	Brazil, French Guiana, Guyana, Suriname

**HONDURAS**

Table 21. The IUCN Mace-Lande Categories for the conservation status of the primates of Honduras.

Species	Category	Distribution
<b>Family Cebidae</b>		
<i>Cebus capucinus limitaneus</i>	LR	Belize, Honduras, Nicaragua
<i>Alouatta palliata palliata</i>	LR	Costa Rica, Nicaragua, Honduras, Panama
<i>Ateles geoffroyi vellerosus</i>	LR	El Salvador, Honduras, Guatemala, Mexico

**MEXICO**

Table 22. The IUCN Mace-Lande Categories for the conservation status of the primates of Mexico.

Species	Category	Distribution
<b>Family Cebidae</b>		
<i>Alouatta palliata mexicana</i>	VU	Guatemala, Mexico
<i>Alouatta pigra</i>	LR	Belize, Guatemala, Mexico
<i>Ateles geoffroyi vellerosus</i>	LR	El Salvador, Honduras, Guatemala, Mexico

**NICARAGUA**

Table 23. The IUCN Mace-Lande Categories for the conservation status of the primates of Nicaragua.

Species	Category	Distribution
<b>Family Cebidae</b>		
<i>Cebus capucinus limitaneus</i>	LR	Belize, Honduras, Nicaragua
<i>Alouatta palliata palliata</i>	LR	Costa Rica, Nicaragua, Honduras, Panama
<i>Ateles geoffroyi frontatus</i>	VU	Costa Rica, Nicaragua

**PANAMA**

Table 24. The IUCN Mace-Lande Categories for the conservation status of the primates of Panama. Endemic taxa are in bold.

Species	Category	Distribution
<b>Family Callitrichidae</b>		
<i>Saguinus geoffroyi</i>	LR	Colombia, Costa Rica, Panama
<b>Family Cebidae</b>		
<i>Aotus lemurinus lemurinus</i>	VU	Colombia, Costa Rica, Panama
<i>Saimiri oerstedi oerstedi</i>	EN	Costa Rica, Panama
<i>Cebus capucinus imitator</i>	LR	Costa Rica, Panama
<i>Alouatta palliata palliata</i>	LR	Costa Rica, Nicaragua, Honduras, Panama
<i>Alouatta palliata aequatorialis</i>	LR	Colombia, Ecuador, Panama, Peru
<i>Alouatta coibensis coibensis</i>	EN	<b>Coiba Island, Panama</b>
<i>Alouatta coibensis trabeata</i>	CR	<b>Panama</b>
<i>Ateles geoffroyi azuerensis</i>	CR	<b>Panama</b>
<i>Ateles geoffroyi grisescens</i>	EN	Colombia, Panama
<i>Ateles geoffroyi panamensis</i>	EN	Costa Rica, Panama
<i>Ateles fusciceps robustus (= rufiventris)</i>	VU	Colombia, Panama

**PARAGUAY**

Table 25. The IUCN Mace-Lande Categories for the conservation status of the primates of Paraguay.

Species	Category	Distribution
<b>Family Callitrichidae</b>		
<i>Callithrix melanura</i>	LR	Bolivia, Brazil, Paraguay
<b>Family Cebidae</b>		
<i>Aotus azarae azarae</i>	LR	Argentina, Bolivia, Paraguay
<i>Callicebus donacophilus pallescens</i>	LR	Bolivia, Brazil, Paraguay
<i>Cebus apella paraguayanus</i>	LR	Argentina, Brazil, Paraguay
<i>Alouatta caraya</i>	LR	Argentina, Brazil, Paraguay, Uruguay (?)

**PERU**

Table 26. The IUCN Mace-Lande Categories for the conservation status of the primates of Peru. Endemic taxa are in bold.

Species	Category	Distribution
<b>Family Callitrichidae</b>		
<i>Cebuella pygmaea</i>	LR	Bolivia, Brazil, Colombia, Ecuador, Peru
<i>Saguinus nigricollis nigricollis</i>	LR	Brazil (?), Colombia, Peru
<i>Saguinus nigricollis graellsi</i>	LR	Colombia, Ecuador, Peru
<i>Saguinus fuscicollis fuscicollis</i>	LR	Brazil, Peru
<i>Saguinus fuscicollis illigeri</i>	<b>LR</b>	<b>Peru</b>
<i>Saguinus fuscicollis leucogenys</i>	<b>LR</b>	<b>Peru</b>
<i>Saguinus fuscicollis nigrifrons</i>	<b>LR</b>	<b>Peru</b>
<i>Saguinus fuscicollis lagonotus</i>	LR	Ecuador, Peru
<i>Saguinus fuscicollis weddelli</i>	LR	Bolivia, Brazil, Peru
<i>Saguinus fuscicollis crandalli</i>	DD	Peru (?)
<i>Saguinus tripartitus</i>	LR	Ecuador, Peru
<i>Saguinus mystax mystax</i>	LR	Brazil, Peru
<i>Saguinus labiatus labiatus</i>	LR	Bolivia, Brazil, Peru
<i>Saguinus imperator imperator</i>	VU	Brazil, Peru
<i>Saguinus imperator subgrisescens</i>	LR	Bolivia, Brazil, Peru
<i>Callimico goeldii</i>	VU	Bolivia, Brazil, Colombia, Peru
<b>Family Cebidae</b>		
<i>Aotus vociferans</i>	LR	Colombia, Brazil, Ecuador, Peru
<i>Aotus miconax</i>	<b>VU</b>	<b>Peru</b>
<i>Aotus nigriceps</i>	LR	Brazil, Colombia, Peru
<i>Aotus azarae boliviensis</i>	LR	Bolivia, Peru
<i>Aotus nancymaae</i>	LR	Brazil, Colombia, Peru
<i>Callicebus oenanthe</i>	<b>VU</b>	<b>Peru</b>
<i>Callicebus brunneus</i>	LR	Brazil, Peru
<i>Callicebus cupreus cupreus</i>	LR	Brazil, Peru
<i>Callicebus cupreus discolor</i>	LR	Colombia, Ecuador, Peru
<i>Callicebus caligatus</i>	LR	Brazil, Peru
<i>Callicebus torquatus lucifer</i>	LR	Brazil, Colombia, Peru
<i>Saimiri boliviensis boliviensis</i>	LR	Bolivia, Brazil, Peru
<i>Saimiri boliviensis peruviensis</i>	<b>LR</b>	<b>Peru</b>
<i>Saimiri sciureus macrodon</i>	LR	Brazil, Colombia, Ecuador, Peru
<i>Cebus apella macrocephalus</i>	LR	Brazil, Peru
<i>Cebus apella maranensis</i>	LR	Brazil, Colombia, Ecuador, Peru
<i>Cebus apella pallidus</i>	LR	Bolivia, Peru
<i>Cebus apella peruanus</i>	LR	Brazil, Peru
<i>Cebus albifrons unicolor</i>	LR	Colombia, Bolivia, Brazil, Peru, Venezuela
<i>Cebus albifrons cuscinus</i>	DD	Bolivia, Brazil, Peru
<i>Cebus albifrons yuracus</i>	DD	Colombia, Ecuador, Peru
<i>Pithecia monachus monachus</i>	LR	Brazil, Colombia, Ecuador, Peru
<i>Pithecia monachus milleri</i>	<b>VU</b>	<b>Colombia</b>
<i>Pithecia irrorata irrorata</i>	LR	Bolivia, Brazil, Peru
<i>Pithecia aequatorialis</i>	LR	Colombia (?), Ecuador, Peru
<i>Cacajao calvus ucayalii</i>	VU	Brazil (?), Peru
<i>Alouatta seniculus seniculus</i>	LR	Colombia, Bolivia, Brazil, Ecuador, Peru, Venezuela
<i>Alouatta seniculus juara</i>	DD	Brazil, Peru
<i>Alouatta palliata aequatorialis</i>	LR	Colombia, Ecuador, Panama, Peru

continued ...

Table 26 (continued)

<i>Ateles belzebuth belzebuth</i>	VU	Brazil, Colombia, Ecuador, Peru, Venezuela
<i>Ateles chamek</i>	LR	Bolivia, Brazil, Peru
<i>Lagothrix lagotricha lagotricha</i>	LR	Brazil, Colombia, Ecuador, Peru
<i>Lagothrix lagotricha cana</i>	VU	Brazil, Peru
<i>Lagothrix lagotricha poeppigii</i>	VU	Brazil, Ecuador, Peru
<i>Lagothrix flavicauda</i>	CR	Peru

**SURINAME**

Table 27. The IUCN Mace-Lande Categories for the conservation status of the primates of Suriname.

Species	Category	Distribution
<b>Family Callitrichidae</b>		
<i>Saguinus midas midas</i>	LR	Brazil, French Guiana, Guyana, Suriname
<b>Family Cebidae</b>		
<i>Saimiri sciureus sciureus</i>	LR	Brazil, French Guiana, Guyana, Suriname
<i>Cebus apella apella</i>	LR	Colombia, Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Cebus olivaceus castaneus</i>	LR	French Guiana, Suriname
<i>Pithecia pithecia pithecia</i>	LR	Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Chiropotes satanas chiropotes</i>	LR	Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Alouatta seniculus macconnelli</i>	LR	Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Ateles paniscus</i>	LR	Brazil, French Guiana, Guyana, Suriname

**TRINIDAD**

Table 28. The IUCN Mace-Lande Categories for the conservation status of the primates of Trinidad. Endemic subspecies are in bold.

Species	Category	Distribution
<b>Family Cebidae</b>		
<i>Cebus albifrons trinitatis</i>	CR	Trinidad
<i>Alouatta seniculus insularis</i>	LR	Trinidad

**URUGUAY**

Table 29. The IUCN Mace-Lande Categories for the conservation status of the primates of Uruguay.

Species	Category	Distribution
<b>Family Cebidae</b>		
<i>Alouatta caraya</i>	LR	Argentina, Brazil, Paraguay, Uruguay (?)

**VENEZUELA**

Table 30. The IUCN Mace-Lande categories for the conservation status of the primates of Venezuela. Endemic taxa are in bold.

Species	Category	Distribution
<b>Family Cebidae</b>		
<i>Aotus trivirgatus</i>	LR	Brazil, Colombia, Venezuela
<i>Callicebus torquatus lugens</i>	LR	Brazil, Colombia, Venezuela
<i>Saimiri sciureus cassiquiarensis</i>	LR	Brazil, Colombia, Venezuela
<i>Cebus apella apella</i>	LR	Colombia, Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Cebus apella margaritae</i>	CR	Venezuela
<i>Cebus albifrons unicolor</i>	LR	Colombia, Bolivia, Brazil, Peru, Venezuela
<i>Cebus albifrons adustus</i>	DD	<b>Venezuela</b>
<i>Cebus albifrons leucocephalus</i>	DD	<b>Venezuela</b>
<i>Cebus olivaceus olivaceus</i>	LR	Brazil, Guyana, Venezuela,
<i>Cebus olivaceus apiculatus</i>	LR	<b>Venezuela</b>
<i>Cebus olivaceus brunneus</i>	LR	<b>Venezuela</b>
<i>Cebus olivaceus</i> ssp.	DD	<b>Venezuela</b>
<i>Pithecia pithecia pithecia</i>	LR	Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Chiropotes satanas chiropotes</i>	LR	Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Cacajao melanocephalus melanocephalus</i>	LR	Brazil, Venezuela
<i>Alouatta seniculus seniculus</i>	LR	Colombia, Bolivia, Brazil, Ecuador, Peru, Venezuela
<i>Alouatta seniculus stramineus</i>	LR	Brazil, Venezuela
<i>Alouatta seniculus macconnelli</i>		Brazil, French Guiana, Guyana, Suriname, Venezuela
<i>Alouatta seniculus</i> ssp.	LR	<b>Venezuela</b>
<i>Alouatta arctoidea</i>	LR	<b>Venezuela</b>
<i>Ateles belzebuth belzebuth</i>	VU	Brazil, Colombia, Ecuador, Peru, Venezuela
<i>Ateles belzebuth hybridus</i>	EN	Colombia, Venezuela
<i>Lagothrix lagotricha lagotricha</i>	LR	Brazil, Colombia, Ecuador, Peru, Venezuela
<i>Lagothrix lagotricha lugens</i>	CR	Colombia, Venezuela

Table 31. A summary of the numbers of Callitrichidae and Cebidae in each of 20 South American countries. Also shown is the number of endemic species and taxa, and the number of threatened taxa in the IUCN Mace Lande categories of Critically Endangered (CR), Endangered (EN) and Vulnerable (VU).

Country	Callitrichidae		Cebidae		Total		Endemic	
	Species	Taxa	Species	Taxa	Species	Taxa	Species	Taxa
<i>Argentina</i>	-	-	4	5	4	5	-	-
<i>Belize</i>	-	-	3	3	3	3	-	-
<i>Bolivia</i>	6	6	11	15	17	21	3	4
<i>Brazil</i>	31	44	45	84	76	128	38	69
<i>Colombia</i>	8	10	23	41	31	51	4	18
<i>Costa Rica</i>	1	1	5	8	6	9	-	2
<i>Ecuador</i>	4	4	12	14	16	18	-	2
<i>El Salvador</i>	-	-	1	1	1	1	-	-
<i>French Guiana</i>	1	1	7	7	8	8	-	-
<i>Guatemala</i>	-	-	3	4	3	4	-	-
<i>Guyana</i>	1	1	7	7	8	8	-	-

continued ...

Table 31 (continued)

<i>Honduras</i>	-	-	3	3	3	3	-	-
<i>Mexico</i>	-	-	3	4	3	4	-	-
<i>Nicaragua</i>	-	-	3	3	3	3	-	-
<i>Panama</i>	1	1	7	11	8	12	1	3
<i>Paraguay</i>	1	1	4	4	5	5	-	-
<i>Peru</i>	8	16	24	35	32	51	3	7
<i>Suriname</i>	1	1	7	7	8	8	-	-
<i>Trinidad</i>	-	-	2	2	2	2	-	2
<i>Uruguay</i>	-	-	1(?)	1(?)	1(?)	-	-	-
<i>Venezuela</i>	-	-	13	24	13	24	1	7

Table 32. The numbers of threatened taxa and threatened endemic taxa in each country according to the Mace-Lande System (IUCN 1995). Critically Endangered (CR), Endangered (EN), and Vulnerable (VU).

Country	Threatened Taxa				Threatened Endemic Taxa			
	CR	EN	VU	Total	CR	EN	VU	Total
<i>Argentina</i>	-	-	1	1	-	-	-	-
<i>Belize</i>	-	-	1	1	-	-	-	-
<i>Bolivia</i>	-	-	1	1	-	-	-	-
<i>Brazil</i>	8	11	19	38	8	10	12	30
<i>Colombia</i>	1	7	10	18	-	4	6	10
<i>Costa Rica</i>	1	2	3	6	1	-	1	2
<i>Ecuador</i>	1	-	2	3	1	-	-	1
<i>El Salvador</i>	-	-	-	-	-	-	-	-
<i>French Guiana</i>	-	-	-	-	-	-	-	-
<i>Guatemala</i>	-	-	2	2	-	-	-	-
<i>Guyana</i>	-	-	-	-	-	-	-	-
<i>Honduras</i>	-	-	-	-	-	-	-	-
<i>Mexico</i>	-	-	2	2	-	-	-	-
<i>Nicaragua</i>	-	-	1	1	-	-	-	-
<i>Panama</i>	2	4	2	8	2	1	-	3
<i>Paraguay</i>	-	-	-	-	-	-	-	-
<i>Peru</i>	1	-	9	10	1	-	2	3
<i>Suriname</i>	-	-	-	-	-	-	-	-
<i>Trinidad</i>	1	-	1	2	1	-	1	2
<i>Uruguay</i>	-	-	-	-	-	-	-	-
<i>Venezuela</i>	2	1	1	4	-	-	-	-

### Regional Primate Diversity

Here we present an analysis of the diversity of primate species and subspecies in seven phytogeographic regions identified by Gentry (1982): Middle America, Northern Andes, Southern Andes, Amazonia (Hylaea), Cerrado/Caatinga/Chaco, and Atlantic Forest (see Figs. 2 and 3, and Tables 33-39).

The analysis of the known or supposed distributions of the 202 primate species and subspecies indicates the following numbers in each of the regions: Middle America - 22; Northern Venezuela-Colombia - 24; Northern Andes - 13; Southern Andes - 6; Amazonia (Hylaea) - 121, Cerrado/Caatinga/Chaco - 11, and Atlantic Forest - 23 (Table .40). Table 41 presents a summary of the number of threatened taxa and the

number of threatened taxa endemic to each region.

From this synthesis it is possible to state the following:

1. There are 22 primates species and subspecies in the Middle American region, 18 of which (82%) are endemic. Thirteen (59%) of the primates occurring in Middle America are threatened. Eleven of the 18 endemic primates (61%) are threatened.
2. There are 24 primates species and subspecies in, 19 of which (79%) are endemic. Ten (42%) of the primates occurring in Northern Venezuela-Colombia are threatened. Eight of the 19 endemic primates (42%) are threatened.

Figure 2. Major phytogeographic regions in the Neotropics [Gentry, 1982] (map by R. B. Machado)



3. There are 13 primate species and subspecies in the Northern Andean region, eight of which (61%) are endemic. Seven (54%) of the primates occurring in the Northern Andes are threatened. Five of the eight endemic primates (62%) are threatened.
4. There are six primate species and subspecies in the Southern Andean region, two of which (33%) are endemic. Three (50%) of the primates occurring in the Southern Andes are threatened. Both of the two endemic primates are threatened.
5. There are 123 primate species and subspecies in the Amazonian region, 112 of which (91%) are endemic. Twenty-two (18%) of the primates occurring in Amazonia are threatened. Twenty-one of the 112 endemic primates (19%) are threatened.

6. There are 11 primate species and subspecies in the Cerrado/Caatinga/Chaco region, five of which (45%) are endemic. None of the primates occurring in the Cerrado/Caatinga/Chaco are considered threatened.
7. There are 23 primate species and subspecies in the Atlantic forest region, 17 of which (74%) are endemic. Eighteen (78%) of the primates occurring in the Atlantic forest are threatened. All of the 18 endemic primates are threatened..



Figure 3. Number of Neotropical primate species and genera in each major phytogeographic region, with number of endemics indicated (map by R. B. Machado).

**MIDDLE AMERICA**

Table 33. Primates occurring in the Middle America phytogeographic region as defined by Gentry (1982). Endemic primates are in bold.

Species	Category	Distribution
<b>Family Callitrichidae</b>		
<i>Saguinus geoffroyi</i>	LR	Middle America, Northern Andes
<b>Family Cebidae</b>		
<i>Aotus lemurinus lemurinus</i>	VU	Middle America, Northern Andes
<i>Saimiri oerstedi oerstedi</i>	EN	Middle America
<i>Saimiri oerstedi citrinellus</i>	CR	Middle America
<i>Cebus capucinus limitaneus</i>	LR	Middle America
<i>Cebus capucinus imitator</i>	LR	Middle America
<i>Alouatta palliata palliata</i>	LR	Middle America
<i>Alouatta palliata mexicana</i>	VU	Middle America
<i>Alouatta palliata aequatorialis</i>	LR	Middle America, Northern Andes
<i>Alouatta coibensis coibensis</i>	EN	Middle America
<i>Alouatta coibensis trabeata</i>	CR	Middle America
<i>Alouatta pigra</i>	LR	Middle America
<i>Ateles geoffroyi geoffroyi</i>	LR	Middle America
<i>Ateles geoffroyi azuerensis</i>	CR	Middle America
<i>Ateles geoffroyi frontatus</i>	VU	Middle America
<i>Ateles geoffroyi grisescens</i>	EN	Middle America
<i>Ateles geoffroyi pan</i>	DD	Middle America
<i>Ateles geoffroyi panamensis</i>	EN	Middle America
<i>Ateles geoffroyi ornatus</i>	VU	Middle America
<i>Ateles geoffroyi vellerosus</i>	LR	Middle America
<i>Ateles geoffroyi yucatanensis</i>	VU	Middle America
<i>Ateles fusciceps robustus</i> (= <i>rufiventris</i> )	VU	Middle America, Northern Venezuela-Colombia

**NORTHERN VENEZUELA-COLOMBIA**

Table 34. Primates occurring in the Northern Venezuela-Colombia phytogeographic region as defined by Gentry (1982). Endemic primates are in bold.

Species	Category	Distribution
<b>Family Callitrichidae</b>		
<i>Saguinus oedipus</i>	EN	Northern Venezuela-Colombia
<b>Family Cebidae</b>		
<i>Aotus lemurinus griseimembra</i>	EN	Northern Venezuela-Colombia
<i>Aotus brumbacki</i>	VU	Northern Venezuela-Colombia
<i>Callicebus cupreus ornatus</i>	VU	Northern-Venezuela-Colombia
<i>Saimiri sciureus albigena</i>	LR	Northern Venezuela-Colombia, Amazonia
<i>Cebus apella margaritae</i>	CR	Northern Venezuela- Colombia
<i>Cebus albifrons albifrons</i>	LR	Northern Venezuela- Colombia
<i>Cebus albifrons adustus</i>	DD	Northern Venezuela- Colombia
<i>Cebus albifrons cesarae</i>	DD	Northern Venezuela- Colombia
<i>Cebus albifrons hypoleucus</i>	DD	Northern Venezuela- Colombia
<i>Cebus albifrons malitiosus</i>	DD	Northern Venezuela- Colombia
<i>Cebus albifrons trinitatis</i>	CR	Northern Venezuela- Colombia
<i>Cebus albifrons leucocephalus</i>	DD	Northern Venezuela- Colombia

continued ...

Table 34 (continued)

<i>Cebus capucinus capucinus</i>	LR	Northern Venezuela- Colombia
<i>Cebus olivaceus apiculatus</i>	LR	Northern Venezuela- Colombia
<i>Cebus olivaceus brunneus</i>	LR	Northern Venezuela- Colombia
<i>Cebus olivaceus</i> ssp.	DD	Northern Venezuela-Colombia
<i>Alouatta seniculus seniculus</i>	LR	Northern Venezuela-Colombia, Amazonia
<i>Alouatta seniculus insulanus</i>	VU	Northern Venezuela-Colombia
<i>Alouatta seniculus</i> ssp.	LR	Northern Venezuela-Colombia
<i>Alouatta arctoidea</i>	LR	Northern Venezuela-Colombia
<i>Ateles fusciceps robustus</i> (= <i>rufiventris</i> )	VU	Northern Venezuela-Colombia, Middle America
<i>Ateles belzebuth hybridus</i>	EN	Northern Venezuela-Colombia, Northern Andes
<i>Ateles belzebuth brunneus</i>	EN	Northern Venezuela-Colombia

**NORTHERN ANDES**

Table 35. Primates occurring in the Northern Andean phytogeographic region as defined by Gentry (1982). Endemic primates are in bold.

Species	Category	Distribution
<b>Family Callitrichidae</b>		
<i>Saguinus leucopus</i>	VU	<b>Northern Andes</b>
<i>Saguinus geoffroyi</i>	LR	Northern Andes, Middle America
<b>Family Cebidae</b>		
<i>Aotus lemurinus lemurinus</i>	VU	Northern Andes, Middle America
<i>Aotus vociferans</i>	LR	Northern Andes, Amazonia
<i>Aotus hershkovitzii</i>	DD	<b>Northern Andes</b>
<i>Callicebus torquatus medemi</i>	VU	<b>Northern Andes</b>
<i>Cebus albifrons aequatorialis</i>	DD	<b>Northern Andes</b>
<i>Cebus albifrons versicolor</i>	DD	<b>Northern Andes</b>
<i>Cebus capucinus curtus</i>	VU	<b>Northern Andes</b>
<i>Alouatta palliata aequatorialis</i>	LR	Northern Andes, Middle America
<i>Ateles fusciceps fusciceps</i>	CR	<b>Northern Andes</b>
<i>Ateles belzebuth hybridus</i>	EN	Northern Andes, Northern Venezuela-Colombia
<i>Lagothrix lagotricha lugens</i>	CR	<b>Northern Andes</b>

**SOUTHERN ANDES**

Table 36. Primates occurring in the Southern Andean phytogeographic region as defined by Gentry (1982). Endemic primates are in bold.

Species	Category	Distribution
<b>Family Cebidae</b>		
<i>Aotus miconax</i>	VU	Southern Andes, Amazonia
<i>Callicebus oenanthe</i>	VU	<b>Southern Andes</b>
<i>Cebus apella maranonis</i>	LR	Southern Andes, Amazonia
<i>Cebus albifrons yuracae</i>	DD	Southern Andes, Amazonia
<i>Alouatta seniculus seniculus</i>	LR	Southern Andes, Northern Venezuela-Colombia, Amazonia
<i>Lagothrix flavicauda</i>	CR	<b>Southern Andes</b>

**AMAZONIA**

Table 37. Primates occurring in the Amazonian phytogeographic region as defined by Gentry (1982). Endemic primates are in bold.

Species	Category	Distribution
<b>Family Callitrichidae</b>		
<i>Cebuella pygmaea</i>	LR	Amazonia
<i>Callithrix argentata</i>	LR	Amazonia
<i>Callithrix leucippe</i>	VU	Amazonia
<i>Callithrix melanura</i>	LR	Amazonia, Chaco
<i>Callithrix intermedia</i>	LR	Amazonia
<i>Callithrix emiliae</i>	LR	Amazonia
<i>Callithrix nigriceps</i>	VU	Amazonia
<i>Callithrix mariae</i>	LR	Amazonia
<i>Callithrix humeralisera</i>	LR	Amazonia
<i>Callithrix chrysoleuca</i>	VU	Amazonia
<i>Callithrix mauesi</i>	LR	Amazonia
<i>Callithrix saterei</i>	DD	Amazonia
<i>Saguinus nigricollis nigricollis</i>	LR	Amazonia
<i>Saguinus nigricollis graellsii</i>	LR	Amazonia
<i>Saguinus nigricollis hernandezi</i>	VU	Amazonia
<i>Saguinus fuscicollis fuscicollis</i>	LR	Amazonia
<i>Saguinus fuscicollis fuscus</i>	LR	Amazonia
<i>Saguinus fuscicollis availapiresi</i>	LR	Amazonia
<i>Saguinus fuscicollis cruzlimai</i>	DD	Amazonia
<i>Saguinus fuscicollis illigeri</i>	LR	Amazonia
<i>Saguinus fuscicollis leucogenys</i>	LR	Amazonia
<i>Saguinus fuscicollis nigrifrons</i>	LR	Amazonia
<i>Saguinus fuscicollis lagonotus</i>	LR	Amazonia
<i>Saguinus fuscicollis weddelli</i>	LR	Amazonia
<i>Saguinus fuscicollis primitivus</i>	LR	Amazonia
<i>Saguinus fuscicollis melanoleucus</i>	LR	Amazonia
<i>Saguinus fuscicollis crandalli</i>	DD	Amazonia
<i>Saguinus tripartitus</i>	LR	Amazonia
<i>Saguinus mystax mystax</i>	LR	Amazonia
<i>Saguinus mystax pileatus</i>	LR	Amazonia
<i>Saguinus mystax pluto</i>	LR	Amazonia
<i>Saguinus labiatus labiatus</i>	LR	Amazonia
<i>Saguinus labiatus thomasi</i>	LR	Amazonia
<i>Saguinus imperator imperator</i>	VU	Amazonia
<i>Saguinus imperator subgrisescens</i>	LR	Amazonia
<i>Saguinus inustus</i>	LR	Amazonia
<i>Saguinus midas midas</i>	LR	Amazonia
<i>Saguinus midas niger</i>	LR	Amazonia
<i>Saguinus bicolor bicolor</i>	EN	Amazonia
<i>Saguinus bicolor ochraceus</i>	LR	Amazonia
<i>Saguinus bicolor martinsi</i>	LR	Amazonia
<i>Callimico goeldii</i>	VU	Amazonia
<b>Family Cebidae</b>		
<i>Aotus vociferans</i>	LR	Amazonia, Northern Andes
<i>Aotus trivirgatus</i>	LR	Amazonia
<i>Aotus miconax</i>	VU	Amazonia, Southern Andes
<i>Aotus nigriceps</i>	LR	Amazonia
<i>Aotus infuscatus</i>	LR	Amazonia, Cerrado
<i>Aotus azarae boliviensis</i>	LR	Amazonia, Chaco

continued ...

Table 37 (continued)

<i>Aotus nancymaae</i>	LR	Amazonia
<i>Callicebus modestus</i>	LR	Amazonia
<i>Callicebus donacophilus donacophilus</i>	LR	Amazonia
<i>Callicebus olallae</i>	DD	Amazonia
<i>Callicebus cinerascens</i>	LR	Amazonia
<i>Callicebus hoffmannsi hoffmannsi</i>	LR	Amazonia
<i>Callicebus hoffmannsi baptista</i>	LR	Amazonia
<i>Callicebus moloch</i>	LR	Amazonia
<i>Callicebus brunneus</i>	LR	Amazonia
<i>Callicebus cupreus cupreus</i>	LR	Amazonia
<i>Callicebus cupreus discolor</i>	LR	Amazonia
<i>Callicebus caligatus</i>	LR	Amazonia
<i>Callicebus dubius</i>	VU	Amazonia
<i>Callicebus torquatus torquatus</i>	LR	Amazonia
<i>Callicebus torquatus lugens</i>	LR	Amazonia
<i>Callicebus torquatus lucifer</i>	LR	Amazonia
<i>Callicebus torquatus purinus</i>	LR	Amazonia
<i>Callicebus torquatus regulus</i>	LR	Amazonia
<i>Saimiri boliviensis boliviensis</i>	LR	Amazonia, Chaco
<i>Saimiri boliviensis peruviensis</i>	LR	Amazonia
<i>Saimiri boliviensis pluvialis</i>	LR	Amazonia
<i>Saimiri boliviensis jaburuensis</i>	LR	Amazonia
<i>Saimiri vanzolinii</i>	VU	Amazonia
<i>Saimiri sciureus sciureus</i>	LR	Amazonia
<i>Saimiri sciureus macrodon</i>	LR	Amazonia
<i>Saimiri sciureus cassiquiarensis</i>	LR	Amazonia
<i>Saimiri sciureus albigena</i>	LR	Amazonia, Northern Venezuela-Colombia
<i>Saimiri ustus</i>	LR	Amazonia
<i>Cebus apella apella</i>	LR	Amazonia
<i>Cebus apella macrocephalus</i>	LR	Amazonia
<i>Cebus apella maranonis</i>	LR	Amazonia, Southern Andes
<i>Cebus apella pallidus</i>	LR	Amazonia
<i>Cebus apella peruanus</i>	LR	Amazonia
<i>Cebus albifrons unicolor</i>	LR	Amazonia
<i>Cebus albifrons cuscinus</i>	DD	Amazonia
<i>Cebus albifrons yuracus</i>	DD	Amazonia, Southern Andes
<i>Cebus olivaceus olivaceus</i>	LR	Amazonia
<i>Cebus olivaceus castaneus</i>	LR	Amazonia
<i>Cebus kaapor</i>	VU	Amazonia
<i>Pithecia pithecia pithecia</i>	LR	Amazonia
<i>Pithecia pithecia chrysocephala</i>	LR	Amazonia
<i>Pithecia monachus monachus</i>	LR	Amazonia
<i>Pithecia monachus milleri</i>	VU	Amazonia
<i>Pithecia irrorata irrorata</i>	LR	Amazonia
<i>Pithecia irrorata vanzolinii</i>	LR	Amazonia
<i>Pithecia albicans</i>	LR	Amazonia
<i>Pithecia aequatorialis</i>	LR	Amazonia
<i>Chiropotes albinasus</i>	LR	Amazonia
<i>Chiropotes satanas satanas</i>	EN	Amazonia
<i>Chiropotes satanas chiropotes</i>	LR	Amazonia
<i>Chiropotes satanas utahicki</i>	VU	Amazonia
<i>Cacajao calvus calvus</i>	EN	Amazonia
<i>Cacajao calvus ucayalii</i>	VU	Amazonia
<i>Cacajao calvus novaesi</i>	EN	Amazonia

continued ...

Table 37 (continued)

<i>Cacajao calvus rubicundus</i>	EN	Amazonia
<i>Cacajao melanocephalus melanocephalus</i>	LR	Amazonia
<i>Cacajao melanocephalus ouakary</i>	LR	Amazonia
<i>Alouatta seniculus seniculus</i>	LR	Amazonia, Northern Venezuela-Colombia
<i>Alouatta seniculus stramineus</i>	LR	Amazonia
<i>Alouatta seniculus macconnelli</i>	LR	Amazonia
<i>Alouatta seniculus amazonica</i>	DD	Amazonia
<i>Alouatta seniculus juara</i>	DD	Amazonia
<i>Alouatta seniculus puruensis</i>	DD	Amazonia
<i>Alouatta sara</i>	LR	Amazonia, Chaco
<i>Alouatta belzebul belzebul</i>	LR	Amazonia
<i>Alouatta belzebul nigerrima</i>	LR	Amazonia
<i>Alouatta belzebul discolor</i>	LR	Amazonia
<i>Alouatta belzebul ululata</i>	CR	Amazonia
<i>Ateles paniscus</i>	LR	Amazonia
<i>Ateles belzebuth belzebuth</i>	VU	Amazonia
<i>Ateles chamek</i>	LR	Amazonia
<i>Ateles marginatus</i>	EN	Amazonia
<i>Lagothrix lagotricha lagotricha</i>	LR	Amazonia
<i>Lagothrix lagotricha cana</i>	VU	Amazonia
<i>Lagothrix lagotricha poeppigii</i>	VU	Amazonia

**CERRADO/CAATINGA/CHACO**

Table 38. Primates occurring in the Cerrado/Caatinga/Chaco phytogeographic region as defined by Gentry (1982). Endemic primates are in bold.

Species	Category	Distribution
<b>Family Callitrichidae</b>		
<i>Callithrix melanura</i>	LR	Chaco, Amazonia
<i>Callithrix jacchus</i>	LR	Caatinga, Atlantic forest
<i>Callithrix penicillata</i>	LR	Cerrado, Atlantic forest
<b>Family Cebidae</b>		
<i>Aotus infuscatus</i>	LR	Cerrado, Amazonia,
<i>Aotus azarai azarai</i>	LR	<b>Chaco</b>
<i>Aotus azarai boliviensis</i>	LR	Chaco, Amazonia
<i>Callicebus donacophilus pallescens</i>	LR	<b>Cerrado/Chaco</b>
<i>Cebus apella libidinosus</i>	LR	<b>Cerrado/Caatinga/Chaco</b>
<i>Cebus apella paraguayanus</i>	LR	<b>Chaco</b>
<i>Alouatta sara</i>	LR	Chaco, Amazonia
<i>Alouatta caraya</i>	LR	<b>Cerrado/Caatinga/Chaco</b>

**ATLANTIC FOREST**

Table 39. Primates occurring in the Atlantic forest phytogeographic region as defined by Gentry (1982). Endemic primates are in bold.

Species	Category	Distribution
<b>Family Callitrichidae</b>		
<i>Callithrix jacchus</i>	LR	Atlantic forest, Caatinga
<i>Callithrix penicillata</i>	LR	Atlantic forest, Cerrado
<i>Callithrix kuhlii</i>	LR	<b>Atlantic forest</b>
<i>Callithrix geoffroyi</i>	VU	<b>Atlantic forest</b>

continued ...

Table 39 (continued)

<i>Callithrix flaviceps</i>	EN	Atlantic forest
<i>Callithrix aurita</i>	EN	Atlantic forest
<i>Leontopithecus rosalia</i>	CR	Atlantic forest
<i>Leontopithecus chrysomelas</i>	EN	Atlantic forest
<i>Leontopithecus chrysopygus</i>	CR	Atlantic forest
<i>Leontopithecus caissara</i>	CR	Atlantic forest
<b>Family Cebidae</b>		
<i>Callicebus personatus personatus</i>	VU	Atlantic forest
<i>Callicebus personatus nigrifrons</i>	VU	Atlantic forest
<i>Callicebus personatus melanochir</i>	VU	Atlantic forest
<i>Callicebus personatus barbarabrowniae</i>	CR	Atlantic forest
<i>Callicebus</i> sp.	CR	Atlantic forest
<i>Cebus apella nigritus</i>	LR	Atlantic forest
<i>Cebus apella robustus</i>	VU	Atlantic forest
<i>Cebus xanthosternos</i>	CR	Atlantic forest
<i>Alouatta belzebul belzebul</i>	LR	Atlantic forest, Amazonia
<i>Alouatta fusca fusca</i>	CR	Atlantic forest
<i>Alouatta fusca clamitans</i>	VU	Atlantic forest
<i>Brachyteles arachnoides</i>	EN	Atlantic forest
<i>Brachyteles hypoxanthus</i>	EN	Atlantic forest

Table 40. A summary of the numbers of Callitrichidae and Cebidae in each of the seven phytogeographic regions as defined by Gentry (1982). Also shown is the number of endemic species and taxa.

Region	Callitrichidae		Cebidae		Total		Endemic Taxa	
	Species	Taxa	Species	Taxa	Species	Taxa	Species	Taxa
Middle America	1	1	7	21	8	22	4	18
Northern Venezuela-Colombia	1	1	12	23	13	24	2	19
Northern Andes	2	2	10	11	12	13	2	8
Southern Andes	-	-	3	6	3	6	2	2
Amazonia	22	42	43	81	65	123	48	112
Cerrado/Caatinga/Chaco	3	3	6	8	9	11	1	5
Atlantic Forest	10	10	8	13	18	23	14	17

Table 41. The numbers of threatened taxa and threatened endemic taxa in each region according to the Mace-Lande System (IUCN 1995). Critically Endangered (CR), Endangered (EN), and Vulnerable (VU).

Region	Threatened Taxa				Threatened Endemic Taxa			
	CR	EN	VU	Total	CR	EN	VU	Total
Middle America	3	4	6	13	3	4	4	11
Northern Venezuela-Colombia	2	4	4	10	2	3	3	8
Northern Andes	2	1	4	7	2	-	3	5
Southern Andes	1	-	2	3	1	-	1	2
Amazonia	-	6	16	22	-	6	15	21
Cerrado/Caatinga/Chaco	-	-	-	-	-	-	-	-
Atlantic Forest	7	5	6	18	7	5	6	18

## References

- Aguirre, A. C. 1971. *O mono Brachyteles arachnoides (É. Geoffroy). Situação atual da Espécie no Brasil.* Academia Brasileira de Ciências, Rio de Janeiro.
- Albúja, L. 1994. Nuevos registros de *Saguinus tripartitus* en la Amazonia ecuatoriana. *Neotropical Primates* 2(2): 8-10.
- Alperin, R. 1993. *Callithrix argentata* (Linnaeus, 1771): considerações taxonômicas e descrição de subespécie nova. *Bol. Mus. Para. Emílio Goeldi, Sér. Zool.* 9(2): 317-328.
- Aquino, R. and Castro N. 1989. Evaluación de *Saguinus labiatus* y otros primates no humanos en Madre de Dios. Informe Anual del Proyecto Peruano de Primatología "Manuel Moro Sommo". 1989. 7pp.
- Aquino, R. and Encarnación, F. 1994. Primates of Peru / Los Primates del Perú. *Primate Report* (40): 1-127.
- Aquino, R. and Encarnación, F. in press. Distribución geográfica de *Saguinus tripartitus* en la Amazonia del Perú. *Neotropical Primates* 3(4).
- Armada, J. L. A., Barroso, C. M. L., Lima, M. M. C., Muniz, J. A. P. C. and Seuánez, H. N. 1987. Chromosome studies in *Alouatta belzebul*. *Am. J. Primatol.* 13: 283-296.
- Ávila-Pires, F. D. de. 1986. On the validity of and geographical distribution of *Callithrix argentata emiliae* Thomas, 1920 (Primates, Callithricidae). In: *A Primatología no Brasil - 2*, M. T. de Mello (ed.), pp.319-322. Sociedade Brasileira de Primatologia, Brasília.
- Ayres, J. M. 1985. On a new species of squirrel monkey, genus *Saimiri*, from Brazilian Amazonia (Primates, Cebidae). *Papéis Avulsos Zoologia*, São Paulo (36): 147-164.
- Ayres, J. M., Mittermeier, R. A. and Constable, I. D. 1982. Brazilian tamarins on the way to extinction? *Oryx* 16: 329-333.
- Barroso, C. M. L. 1995. Filogenia molecular da subfamília Callitrichinae (*sensu* Rosenberger, 1981). Doctoral Thesis, Universidade Federal do Pará, Belém.
- Bender, M. A. and Chu, E. H. Y. 1963. The chromosomes of primates. In: *Evolutionary Biology and Genetic Biology of the Primates*, J. Buettner-Janusch (ed.), pp.261-310. Academic Press, New York.
- Bicca-Marques, J. C. and Calegaro-Marques, C. 1995. Updating the known distribution of the pygmy marmoset (*Cebuella pygmaea*) in the state of Acre, Brazil. *Neotropical Primates* 3(2): 48-49.
- Bitteti, M. S. di, Placci, G., Brown, D. and Rode, D. I. 1994. Conservation and population status of the brown-howling monkey (*Alouatta fusca clamitans*) in Argentina. *Neotropical Primates* 2(4): 1-4.
- Bodini, R. 1989. Distribución geográfica y conservación de primates no humanos em Colombia. In: *La Primatología en Latinoamérica*, C. J. Saavedra, R. A. Mittermeier and I. B. Santos (eds.), pp.101-113. World Wildlife Fund - U.S., Washington, D. C.
- Bodini, R. and Pérez-Hernández, R. 1987. Distribution of the species and subspecies of cebids in Venezuela. *Fieldiana, Zoology, New Series*, (39): 231-244.
- Bonvicino, C. R., Langguth, A., and Mittermeier, R. A. 1989. A study of the pelage color and geographic distribution in *Alouatta belzebul* (Primates: Cebidae). *Rev. Nordestina Biol.* 6(2): 139-148.
- Boubli, J. P. 1994. The black uakari monkey in the Pico da Neblina National Park. *Neotropical Primates* 2(3): 11-12.
- Brown, A. D. and Rumiz, D. I. 1986. Distribución de los primates en Bolivia. In: *A Primatología no Brasil - 2*, M. T. de Mello (ed.), pp.335-363. Sociedade Brasileira de Primatologia, Brasília.
- Butchart, S. H. M., Barnes, R., Davies, C. W. N., Fernandez, M. and Seddon, N. In press. Observations of two threatened primates in the Peruvian Andes. *Primate Conservation* (16).
- Cabrera, A. 1957. Catalogo de los mamíferos de América del Sur. *Rev. Mus. Argentino de Cienc. Nat. "Bernardino Rivadavia"* 4(1): 1-307.
- Cheverud, J. M. and Moore, A. J. 1990. Subspecific variation in the saddle-back tamarin (*Saguinus fuscicollis*). *Am. J. Primatol.* 21: 1-15.
- Chu, E. H. Y. and Bender, M. A. 1961. Chromosome cytology and evolution in primates. *Science* 133: 1399-1405.
- Coimbra-Filho, A. F. 1969. Mico-leão, *Leontideus rosalia* (Linnaeus, 1766), situação atual da espécie no Brasil. *An. Acad. Brasil. Ciênc.* 41 (supl.): 29-52.
- Coimbra-Filho, A. F. 1970. Acerca de um caso de hibridismo entre *Callithrix jacchus* (L. 1758) x *Callithrix geoffroyi* (Humboldt, 1812). *Rev. Brasil. Biol.* 30: 507-514.
- Coimbra-Filho, A. F. 1971. Os sagüis do gênero *Callithrix* da região oriental brasileira e um caso de duplo-hibridismo entre três de suas formas (Callithricidae, Primates). *Rev. Brasil. Biol.* 31:377-388.
- Coimbra-Filho, A. F. 1972. Mamíferos ameaçados de extinção no Brasil. In: *Espécies da Fauna Brasileira Ameaçadas de Extinção*, pp.13-98. Academia Brasileira de Ciências, Rio de Janeiro.
- Coimbra-Filho, A. F. 1976. *Leontopithecus rosalia chrysopygus* (Mikan, 1823), o mico-leão do Estado de São Paulo (Callitrichidae, Primates). *Silvic. S. Paulo* 10: 1-36.
- Coimbra-Filho, A. F. 1982. Distribuição geográfica, ecologia, extinção e preservação dos platirrinos. In: *Genética Comparada de Primatas Brasileiros*, P. H. Saldaña (ed.), pp.83-103. Sociedade Brasileira de Genética, São Paulo.
- Coimbra-Filho, A. F. 1984. Situação atual dos calitriquídeos que ocorrem no Brasil (Callitrichidae-Primates). In: *A Primatología no Brasil*, M.T.de Mello (ed.), pp.15-33. Sociedade Brasileira de Primatologia, Brasília.
- Coimbra-Filho, A. F. 1985. Sagüi-de-Wied *Callithrix kuhli* (Weid, 1826). *FBCN/Inf.*, Rio de Janeiro 9(4): 5.
- Coimbra-Filho, A. F. 1986a. Sagüi-da-serra *Callithrix flaviceps* (Thomas, 1903). *FBCN/Inf.*, Rio de Janeiro 10(1): 3.
- Coimbra-Filho, A. F. 1986b. Sagüi-da-serra-escuro *Callithrix aurita* (É. Geoffroy, 1812). *FBCN/Inf.*, Rio de Janeiro 10(2): 3.
- Coimbra-Filho, A. F. 1987. Sagüis-de-cara-nua *Saguinus bicolor* (Spix, 1823). *FBCN/Inf.*, Rio de Janeiro 11(1): 3.
- Coimbra-Filho, A. F. 1990. Sistemática, distribuição geográfica e situação atual dos símios brasileiros (Platyrrhini - Primates). *Rev. Brasil. Biol.*, 50: 1063-1079.

- Coimbra-Filho, A. F. 1991. Apontamentos sobre *Callithrix aurita* (É. Geoffroy, 1812), um sagüi pouco conhecido. In: *A Primatologia no Brasil - 3*, A. B. Rylands, and A. T. Bernardes (eds.), pp. 145-158. Sociedade Brasileira de Primatologia and Fundação Biodiversitas, Belo Horizonte.
- Coimbra-Filho, A. F. and Mittermeier, R. A. 1972. Taxonomy of the genus *Leontopithecus* Lesson, 1840. In: *Saving the Lion Marmoset*, D. D. Bridgwater (ed.), pp. 7-22. Wild Animal Propagation Trust, Wheeling, West Virginia.
- Coimbra-Filho, A. F. and Mittermeier, R. A. 1973a. Distribution and ecology of the genus *Leontopithecus* Lesson, 1840 in Brazil. *Primates* 14(1): 47-66.
- Coimbra-Filho, A. F. and Mittermeier, R. A. 1973b. New data on the taxonomy of the Brazilian marmosets of the genus *Callithrix* Erxleben 1777. *Folia Primatol.* 20: 241-264.
- Coimbra-Filho, A. F. and Mittermeier, A. A. 1977. Conservation of the Brazilian lion tamarins (*Leontopithecus rosalia*). In: *Primate Conservation*, H. S. H. Prince Rainier III of Monaco and G. H. Bourne (eds.), pp. 59-94. Academic Press, New York.
- Coimbra-Filho, A. F., Rocha e Silva, R. da and Pissinatti, A. 1991. Acerca da distribuição geográfica original de *Cebus apella xanthosternos* Wied, 1820 (Cebidae, Primates). In: *A Primatologia no Brasil - 3*, A. B. Rylands and A. T. Bernardes (eds.), pp. 215-224. Fundação Biodiversitas and Sociedade Brasileira de Primatologia, Belo Horizonte.
- Coimbra-Filho, A. F., Pissinatti, A. and Rylands, A. B. 1993. Breeding muriquis (*Brachyteles arachnoides*) in captivity: the experience of the Rio de Janeiro Primate Centre (CPRJ-FEEMA) (Ceboidea, Primates). *Dodo, J Wildl. Preserv. Trusts*, 29: 66-77.
- Conservation International. 1990. *The Rain Forest Imperative: A Ten Year Strategy to Save Earth's Most Threatened Ecosystems*. Conservation International, Washington, D. C. 15pp.
- Costello, R. K., Dickinson, C., Rosenberger, A. L., Boinski, S. and Szalay, F. S. 1993. Squirrel monkey (genus *Saimiri*) taxonomy: a multidisciplinary study of the biology of the species. In: *Species, Species Concepts, and Primate Evolution*, W. H. Kimbel and L. B. Martin (eds.), pp. 177-210. Plenum Press, New York.
- Curdts, T. 1993. Distribución geográfica de las dos especies de mono zaraguate que habitan en Guatemala, *Alouatta palliata* y *Alouatta pigra*. In: *Estudios Primatólogicas en México*, Vol. 1, A. Estrada, E. Rodríguez-Luna, R. Lopes-Wilchis and R. Coates-Estrada (eds.), pp. 317-329. Universidad Veracruzana, Veracruz.
- Dahl, J. F. 1987. Conservation of primates in Belize, Central America. *Primate Conservation* (8): 119-121.
- De Boer, L. E. M. 1974. Cytotaxonomy of the Platyrhini (Primates). *Genen. Phaenen.* 17: 1-115.
- De Boer, L. E. M. and De Brujin, M. 1990. Chromosomal distinction between the red-faced and black-faced black spider monkeys (*Ateles paniscus paniscus* and *A. p. chamek*). *Zoo Biol.* 9: 307-316.
- Defler, T. R. 1989. The status and some ecology of primates in the Colombian Amazon. *Primate Conservation* (10): 51-56.
- Defler, T. R. 1994. La conservación de primates en Colombia. *Trianea (Act. Cien. INDERENA)* 5: 255-287.
- Della Serra, O. 1951. Divisão do gênero *Leontocebus* (Macacos, Platyrhini) em dois subgêneros sob bases de caracteres dento-morfológicos. *Pap. Avulsos São Paulo*, 10(8): 147-154.
- Dollman, G. 1933. *Primates. Ser. 3*, British Museum (Natural History), London.
- Dutrillaux, B. 1988. New interpretation of the presumed common ancestral karyotype of platyrhine monkeys. *Folia Primatol.* 50: 226-229.
- Egler, S. G. 1983. Current status of the pied bare-face tamarin in Brazilian Amazonia. *IUCN/SSC Primate Specialist Group Newsletter* (3): 20.
- Eisenberg, J. F. 1989. *Mammals of the Neotropics: The Northern Neotropics Vol. 1, Panama, Colombia, Venezuela, Guyana, Suriname, French Guiana*. Chicago University Press, Chicago.
- Elliot, D. G. 1913. *A Review of Primates*, Monograph Series, American Museum of Natural History, New York.
- Encarnación, F. 1994. Revision of 1990 IUCN Red List of Threatened Animals, IUCN SSC Primate Specialist Group. Primer Borrador para los primates del Perú. Unpublished report to IUCN/SSC Primate Specialist Group (PSG), July 1993. pp. 34.
- Fernandes, M. E. B. 1993. New field records of night monkeys, genus *Aotus*, in northern Brazil. *Neotropical Primates* 1(4): 6-8.
- Ferrari, S. F. 1993. An update on the black-headed marmoset, *Callithrix nigriceps* Ferrari and Lopes 1992. *Neotropical Primates* 1(4): 11-13.
- Ferrari, S. F. 1994. The distribution of the black-headed marmoset, *Callithrix nigriceps*: a correction. *Neotropical Primates* 2(1): 11-12.
- Ferrari, S. F. 1995. Observations on *Chiropotes albinasus* from the Rio dos Marmelos, Amazonas, Brazil. *Primates* 36(2): 289-293.
- Ferrari, S. F. and Lopes, M. A. 1992. A new species of marmoset, genus *Callithrix* Erxleben 1777 (Callitrichidae, Primates) from western Brazilian Amazonia. *Goeldiana Zoologia* (12), 1-3.
- Fonseca, G. A. da, Lemos de Sá, R. M., Pope, T. R., Glander, K. E. and Struhsaker, T. T. 1991. A pilot study of genetic and morphological variation in the muriqui (*Brachyteles arachnoides*) as a contribution to a long-term conservation management plan. Unpublished report to World Wildlife Fund - U. S., Washington, D. C. 58pp.
- Fonseca, G. A. B. da, Cavalcanti, R., Santos, I. B. and Braga, R. 1995. Priority areas for conservation in the Atlantic forest of north-east Brazil. *Neotropical Primates* 3(2): 54-55.
- Fooden, J. 1963. A revision of the woolly monkeys (genus *Lagothrix*). *J. Mammal.* 44(2): 213-247.
- Forman, L., Kleiman, D. G., Bush, R. M., Dietz, J. M., Ballou, J. D., Phillips, L. G., Coimbra-Filho, A. F. and O'Brien, S. J. 1986. Genetic variation within and among lion tamarins. *Am. J. Phys. Anthropol.* 71: 1-11.
- Froehlich, J. W. and Froehlich, P. H. 1986. Dermatoglyphics and subspecific systematics of mantled howler monkeys (*Alouatta palliata*). In: *Current Perspectives in Pri-*

- mate Biology*, D. M. Taub and F. A. King (eds.), pp. 107-121. Van Nostrand Reinhold, New York.
- Froehlich, J. W. and Froehlich, P. H. 1987. The status of Panama's endemic howling monkeys. *Primate Conservation* (8): 58-62.
- Froehlich, J. W., Supriatana, J. and Froehlich, P. H. 1991. Morphometric analyses of *Ateles*: systematic and biogeographic implications. *Am. J. Primatol.* 25: 1-22.
- Galbreath, G. J. 1983. Karyotypic evolution in *Aotus*. *Am. J. Primatol.* 4: 245-251.
- Gentry, A. 1982. Neotropical floristic diversity: phytogeographical connections between Central and South America, Pleistocene climatic fluctuations, or an accident of the Andean orogeny? *Ann. Missouri Bot. Gard.* 69: 557-593.
- Groombridge, B. 1993. *1994 IUCN Red List of Threatened Animals*. World Conservation Monitoring Center, Cambridge, UK.
- Groves, C. P. 1989. *A Theory of Human and Primate Evolution*. Oxford University Press, New York.
- Groves, C. P. 1992. Titis, New World monkeys of the genus *Callicebus* (Cebidae, Platyrhini): a preliminary taxonomic review. *Fieldiana, Zoology, New Series*, (55): 1-109.
- Hershkovitz, P. 1990. *Int. J. Primatol.* 13: 111-112. (Review).
- Groves, C. P. 1993. Order Primates. In: *Mammal Species of the World: A Taxonomic and Geographic Reference*, 2nd Edition, D. E. Wilson and D. M. Reeder (eds.), pp. 243-277. Smithsonian Institution Press, Washington, D. C.
- Hanihari T. and Natori, M. 1987. Preliminary analysis of numerical taxonomy of the genus *Saguinus* based on dental measurements. *Primates* 28(4): 517-523.
- Heltne, P. G. and Kunkel L. M. 1975. Taxonomic notes on the pelage of *Ateles paniscus paniscus*, *A. p. chamek* (sensu Kellogg and Goldman, 1944) and *A. fusciceps rufiventris* (= *A. f. robustus*, Kellogg and Goldman, 1944). *J. Med. Primatol.* 4: 83-102.
- Hernández-Camacho, J. and Cooper, R. W. 1976. The non-human primates of Colombia. In: *Neotropical Primates: Field Studies and Conservation*, R. W. Thorington Jr. and P. G. Heltne (eds.), pp. 35-69. National Academy of Sciences, Washington, D. C.
- Hernández-Camacho, J. and Defler, T. R. 1988. Some aspects of the conservation of non-human primates in Colombia. *Primate Conservation* (6): 42-50.
- Hernández-Camacho, J. and Defler, T. R. 1991. Algunos aspectos de la conservación de primates no-humanos en Colombia. In: *La Primatología en Latinoamérica*, C. J. Saavedra, R. A. Mittermeier and I. B. Santos (eds.), pp. 67-100. World Wildlife Fund, Washington, D.C.
- Hershkovitz, P. 1949. Mammals of northern Colombia. Preliminary report No. 4: Monkeys (Primates), with taxonomic revisions of some forms. *Proc. U. S. Nat. Mus.* 98: 323-427.
- Hershkovitz, P. 1972. Notes on New World Monkeys. *Int. Zoo Yearbook* 12: 3-12.
- Hershkovitz, P. 1977. *Living New World Monkeys (Platyrrhini) with an Introduction to Primates*, Vol. 1. Chicago University Press, Chicago.
- Hershkovitz, P. 1979. Races of the emperor tamarin, *Saguinus imperator* Goeldi (Callitrichidae, Primates). *Primates* 20(2): 277-287.
- Hershkovitz, P. 1982. Subspecies and geographic distribution of black-mantle tamarins *Saguinus nigricollis* Spix (Primates: Callitrichidae). *Proc. Biol. Soc. Wash.* 95(4): 647-656.
- Hershkovitz, P. 1983. Two new species of night monkeys, genus *Aotus* (Cebidae, Platyrhini): a preliminary report on *Aotus* taxonomy. *Am. J. Primatol.* 4(3): 209-243.
- Hershkovitz, P. 1984. Taxonomy of squirrel monkeys, genus *Saimiri*, (Cebidae, Platyrhini): a preliminary report with description of a hitherto unnamed form. *Am. J. Primatol.* 4: 209-243.
- Hershkovitz, P. 1985. A preliminary taxonomic review of the South American bearded saki monkeys genus *Chiropotes* (Cebidae, Platyrhini), with the description of a new subspecies. *Fieldiana, Zoology, New Series* (27): iii + 46.
- Hershkovitz, P. 1987a. The taxonomy of South American sakis, genus *Pithecia* (Cebidae, Platyrhini): a preliminary report and critical review with the description of a new species and new subspecies. *Am. J. Primatol.* 12: 387-468.
- Hershkovitz, P. 1987b. Uacaries, New World monkeys of the genus *Cacajao* (Cebidae, Platyrhini): a preliminary taxonomic review with the description of a new subspecies. *Am. J. Primatol.* 12: 1-53.
- Hershkovitz, P. 1990. Titis, New World monkeys of the genus *Callicebus* (Cebidae, Platyrhini): a preliminary taxonomic review. *Fieldiana, Zoology, New Series*, (55): 1-109.
- Hill, W. C. O. 1957. *Primates. Comparative Anatomy and Taxonomy III. Hapalidae*. Edinburgh University Press, Edinburgh.
- Hill, W. C. O. 1960. *Primates. Comparative Anatomy and Taxonomy IV. Cebidae Part A*. Edinburgh University Press, Edinburgh.
- Hill, W. C. O. 1962. *Primates. Comparative Anatomy and Taxonomy V. Cebidae Part B*. Edinburgh University Press, Edinburgh.
- Hirsch, A., Landau, E. C., Tedeschi, A. C. de M., Meneghetti, J. O. 1991. Estudo comparativo das espécies do gênero *Alouatta* Lacépède, 1799 (Platyrrhini, Atelidae) e sua distribuição geográfica na América do Sul. In: *A Primatologia no Brasil - 3*, A. B. Rylands and A. T. Bernardes (eds.), pp. 239-262. Fundação Biodiversitas and Sociedade Brasileira de Primatologia, Belo Horizonte.
- Horwich, R. H. 1983. Species status of the black howler monkey, *Alouatta pigra*, of Belize. *Primates* 24: 288-289.
- Horwich, R. H. and Johnson, E. D. 1984. Geographic distribution and status of the black howler monkey. *IUCN/SSC Primate Specialist Group Newsletter* (4): 25-27.
- Husson, A. M. 1978. *The Mammals of Suriname*. E. J. Brill, Leiden.
- IUCN. 1993. *Draft IUCN Red List Categories*. The World Conservation Union (IUCN), Species Survival Commission, Gland.
- IUCN. 1995. *IUCN Red List Categories*. The World Conservation Union (IUCN), Species Survival Commission, Gland.

- Kellogg R. and Goldman, E. A. 1944. Review of the spider monkeys. *Proc. U. S. Nat. Mus.* 96: 1-45.
- Kierulff, M. C. M. 1993. Status and distribution of the golden lion tamarin in Rio de Janeiro. *Neotropical Primates* 1(4): 23-24.
- Kierulff, M. C. M. and Oliveira, P. P. de. 1994. Habitat preservation and the translocation of threatened groups of golden lion tamarins, *Leontopithecus rosalia*. *Neotropical Primates* 3(suppl.): 15-18.
- Kinzey, W. G. 1982. Distribution of primates and forest refuges. In: *Biological diversification in the tropics*, G. T. Prance (ed.), pp.455-482. Columbia University Press, New York.
- Kobayashi, S. 1995. A phylogenetic study of titi monkeys, genus *Callicebus*, based on cranial measurements: I. Phyletic groups of *Callicebus*. *Primates* 36(1): 101-120.
- Kobayashi, S. and Langguth, A. 1994a. New titi monkey from Brazil. *Reichurui Kenkyu / Primate Research* 10(2): 128.
- Kobayashi, S. and Langguth, A. L. 1994b. New titi monkey from Brazil. In: *Handbook and Abstracts, XV Congress of the International Primatological Society*, p.166. Kuta, Bali, Indonesia, 3-8 August, 1994.
- Konstant, W., Mittermeier, R. A. and Nash, S. D. 1985. Spider monkeys in captivity and in the wild. *Primate Conservation* (5): 82-109.
- Kunkel, L. M., Heltne, P. G. and Borgaonkar, D. S. 1980. Chromosomal variation and zoogeography in *Ateles*. *Int. J. Primatol.* 1(3): 223-232.
- Langguth, A., Teixeira, D. M., Mittermeier, R. A. and Bonvicino, C. 1987. The red-handed howler monkey in northeastern Brazil. *Primate Conservation* (8): 36-39.
- Lemos de Sá, R. M. and Glander, K. E. 1993. Capture techniques and morphometrics for the woolly spider monkey, or muriqui (*Brachyteles arachnoides*, E. Geoffroy 1806). *Am. J. Primatol.* 29: 145-153.
- Lemos de Sá, R. M., Pope, T. R., Glander, K. E., Struhsaker, T. T. and Fonseca, G. A. B. da 1990. A pilot study of genetic and morphological variation in the muriqui (*Brachyteles arachnoides*). *Primate Conservation* (11): 26-30.
- Lemos de Sá, R. M., Pope, T. R., Struhsaker, T. T. and Glander, K. E. 1993. Sexual dimorphism in canine length of woolly spider monkeys (*Brachyteles arachnoides*, E. Geoffroy 1806). *Int. J. Primatol.* 14(5): 755-763.
- Leo Luna, M. 1987. Primate conservation in Peru: a case study of the yellow-tailed woolly monkey. *Primate Conservation* (8): 122-123.
- Lima, M. M. C. and Seuánez, H. N. 1989. Cytogenetic characterization of *Alouatta belzebul* with atypical pelage coloration. *Folia Primatol.* 52: 97-101.
- Lima, M. M. C. and Seuánez, H. N. 1991. Chromosome studies in the red howler monkey, *Alouatta seniculus stramineus* (Platyrrhini, Primates): description of an  $X_1X_2Y_1Y_2/X_1X_1X_2X_2$  sex-chromosome system and karyological comparison with other subspecies. *Cytogen. Cell Genet.* 57: 151-156.
- Lima, M. M. C., Sampaio, M. I. C., Schneider, M. P. C., Scheffrahn, W., Schneider, H. and Salzano, F. M. 1990. Chromosome and protein variation in red howler monkeys. *Brazil. J. Genet.* 13(4): 789-802.
- Lorini, V. G. and M. L. Persson. 1990. Uma nova espécie de *Leontopithecus* Lesson, 1840, do sul do Brasil (Primates, Callitrichidae). *Bol. Mus. Nacional, Rio de Janeiro, nova sér. Zoologia*,338:1-14.
- Lorini, M. L. and Persson, V. G. 1994. Status of field research on *Leontopithecus caissara*: the Black-Faced Lion Tamarin Project. *Neotropical Primates* 2(suppl.): 52-55.
- Mace, G. M. 1994a. An investigation into methods for categorizing the conservation status of species. In: *Large-Scale Ecology and Conservation Biology*, P. J. Edwards, R. M. May and N. R. Webb, pp.293-312. Blackwell Scientific Publications, Oxford.
- Mace, G. M. 1994b. Classifying threatened species: means and ends. *Phil. Trans. Roy. Soc., Lond.* B344: 91-97.
- Mace, G. M. 1995. Classification of threatened species and its role in conservation planning. In: *Extinction Rates*, J. H. Lawton and R. M. May (eds.), pp.197-213. Oxford University Press, Oxford.
- Mace, G. M. and Lande, R. 1991. Assessing extinction threats: toward a re-evaluation of IUCN threatened species categories. *Conservation Biology* 5: 148-157.
- Mace, G. M.. and Stuart, S. N. 1994. Draft IUCN Red List Categories, Version 2.2. *Species* (21-22): 13-24.
- Mace, G. M., Collar, N., Cooke, J., Gaston, K., Ginsberg, J., Leader-Williams, N., Maunder, M. and Milner-Gulland, E. J. 1992. The development of new criteria for listing species on the IUCN Red List. *Species* (19): 16-22.
- Marroig, G. 1995. Espécies ou subespécies em *Callithrix*? *Neotropical Primates* 3(1): 10-13.
- Martin, R. D. 1992. Goeldi and the dwarfs: the evolutionary biology of the small New World monkeys. *J. Hum. Evol.*, 22: 367-393.
- Martins, E. S., Ayres, J. M. and do Valle, M. B. R. 1988. On the status of *Ateles belzebuth marginatus* with notes on other primates of the Iriri river basin. *Primate Conservation* (9): 87-91.
- Martuscelli, P., Petroni, L. M. and Olmos, F. 1994. Fourteen new localities for the muriqui *Brachyteles arachnoides*. *Neotropical Primates*, 2(2): 12-15.
- Mast, R. B., Rodríguez, J. V. and Mittermeier, R. A. 1993. The Colombian cotton-top tamarin in the wild. In: *A Primate Model for the Study of Colitis and Colonic Carcinoma: the Cotton-top Tamarin* (*Saguinus oedipus*), N. K. Clapp (ed.), pp.3-43. CRC Press, Inc., Oak Ridge.
- McNeely, J. A., Miller, K. R., Reid, W. V., Mittermeier, R. A. and Werner, T. B. 1990. *Conserving the World's Biological Diversity*. International Union for Conservation of Nature and Natural Resources (IUCN), Gland.
- Medeiros, M. A. A. 1994. Citogenética, Evolução Cromossômica, Radiação e Especiação dos Macacos-Aranha (*Ateles*, Primates). Unpubl. Master's Thesis, Universidade Federal do Pará, Museu Paraense Emílio Goeldi, Belém, Pará. 137pp.
- Meireles, C. M. M., Sampaio, N. I. C., Schneider, H. and Schneider, M. P. C. 1992. Protein variation, taxonomy and differentiation in five species of marmosets (genus *Callithrix* Erxleben, 1777). *Primates* 33(2): 227-238.
- Melo, A. C. A., Sampaio, M. I. C., Schneider, M. P. C. and Schneider, H. 1992. Biochemical diversity and genetic distance in two species of the genus *Saguinus*. *Primates*

- 33(2): 217-225.
- Mendes, S. L. 1993. Distribuição geográfica e estado de conservação de *Callithrix flaviceps* (Primates, Callitrichidae). In: *A Primatologia no Brasil - 4*, M. E. Yamamoto and M. B. C. de Souza, pp.139-154. Sociedade Brasileira de Primatologia, Natal.
- Minezawa, M., Harada, M., Jordan, O. C. and Valdivia Borda, C. J. 1985. Cytogenetics of the Bolivian endemic red howler monkeys (*Alouatta seniculus sara*): accessory chromosomes and Y-autosome translocation related numerical variations. *Kyoto University Overseas Research Reports of New World Monkeys* 5: 7-16.
- Mittermeier, R. A. 1987. Framework for primate conservation in the Neotropical region. In: *Primate Conservation in the Tropical Rain Forest*, C. W. Marsh and R. A. Mittermeier (eds.), pp.305-320. Alan R. Liss, New York.
- Mittermeier, R. A. and Coimbra-Filho, A. F. 1981. Systematics: species and subspecies. In: *Ecology and Behavior of Neotropical Primates*, Vol. 1, A. F. Coimbra-Filho and R. A. Mittermeier (eds.), pp.29-109. Academia Brasileira de Ciências, Rio de Janeiro.
- Mittermeier, R. A. and Oates, J. F. 1985. Primate diversity: the World's top countries. *Primate Conservation* (5): 41-48.
- Mittermeier, R. A., Valle, C. M. C., Alves M. C., Santos, I. B., Pinto, C. A. M., Strier, K. B., Young, A. L., Veado, E. M., Constable, I. D., Paccagnella, S. G., and Lemos de Sá, R. M. 1987. Current distribution of the muriqui in the Atlantic forest region of eastern Brasil. *Primate Conservation* (8):143-148.
- Mittermeier, R. A., Rylands, A. B. and Coimbra-Filho, A. F. 1988. Systematics: species and subspecies - an update. In: *Ecology and Behavior of Neotropical Primates*, Vol. 2, R. A. Mittermeier, A. B. Rylands, A. F. Coimbra-Filho and G. A. B. da Fonseca (eds.), pp.13-75. World Wildlife Fund, Washington, D.C.
- Mittermeier, R. A., Schwarz, M., Ayres, J. M. 1992a. A new species of marmoset, genus *Callithrix* Erxleben 1777 (Callitrichidae, Primates), from the Rio Maués region, state of Amazonas, Central Brazilian Amazonia. *Goeldiana Zoologia* (14): 1-17.
- Mittermeier, R. A., Werner, T., Ayres, J. M. and Fonseca, G. A. B. da. 1992b. O país da megadiversidade. *Ciência Hoje*, 14(81): 20-27.
- Moore, A. J. and Cheverud, J. M. 1992. Systematics of the *Saguinus oedipus* group of the bare-faced tamarins: evidence from facial morphology. *Am. J. Phys. Anthropol.* 89: 73-84.
- Muskin, A. 1984. Field notes and geographic distribution of *Callithrix aurita* in eastern Brazil. *Am. J. Primatol.* 7: 377-380.
- Myers, N. 1987. Trends in the destruction of rain forests. In: *Primate Conservation in the Tropical Rain Forest*, C. W. Marsh and R. A. Mittermeier (eds.), pp.3-22. Alan R. Liss, New York.
- Myers, N. 1988. Threatened biotas: "hot spots" in tropical forests. *The Environmentalist* 8(3): 187-208.
- Myers, N. 1990. The biodiversity challenge: expanded hot-spots analysis. *The Environmentalist* 10(4): 243-265.
- Napier, J. S. and Napier, P. H. 1967. *A Handbook of Living Primates*. Academic Press, London.
- Natori, M. 1986. Interspecific relationships of *Callithrix* based on dental characters. *Primates* 27(3): 321-336.
- Natori, M. 1989. An analysis of cladistic relationships of *Leontopithecus* based on dental and cranial characters. *J. Anthropol. Soc. Nippon* 97(2): 157-167.
- Natori, M. 1990. Numerical analysis of the taxonomical status of *Callithrix kuhli* based on the measurements of the postcanine dentition. *Primates* 31(4): 555-562.
- Natori, M. 1994. Craniometrical variations among eastern Brazilian marmosets and their systematic relationships. *Primates* 35(2): 167-176.
- Natori, M. and Hanihara, T. 1992. Variations in dental measurements between *Saguinus* species and their systematic relationships. *Folia Primatol.* 58: 84-92.
- Natori, M. and Shigehara, N. 1992. Interspecific differences in lower dentition among eastern Brazilian marmosets. *J. Mammal.* 73(3): 668-671.
- Oliver, W. L. R. and I. B. Santos. 1991. Threatened endemic mammals of the Atlantic forest region of southeast Brazil. *Wildl. Preserv. Trust, Special Scientific Report* (4): 1-125.
- Olmos, F. and Martuscelli, P. 1995. Habitat and distribution of the buffy-tufted-ear marmoset *Callithrix aurita* in São Paulo State, Brazil, with notes on its natural history. *Neotropical Primates* 3(3): 75-79.
- Peres, C. A. 1993. Notes on the primates of the Juruá river, western Brazilian Amazonia. *Folia Primatol.* 61: 97-103.
- Pieczarka, J. C. and Nagamachi, C. Y. 1988. Cytogenetic studies of *Aotus* from eastern Amazonia: Y/autosome rearrangement. *Am. J. Primatol.* 14: 255-263.
- Pieczarka, J. C., Barros, R. M. de S., Nagamachi, C. Y., Rodrigues, R. and Espinel, A. 1992. *Aotus vociferans* x *Aotus nancymai*: sympatry without chromosomal hybridation. *Primates* 33(2): 239-245.
- Pinto, L. P. S. and Tavares, L. I. 1994. Inventory and conservation status of wild populations of golden-headed lion tamarins, *Leontopithecus chrysomelas*. *Neotropical Primates* 2(suppl.): 24-27.
- Pocock, R. I. 1925. Additional notes on the external characters of some platyrhine monkeys. *Proc. Zool. Soc. Lond.* (1925): 27-47.
- Queiroz, H. L. 1992. A new species of capuchin monkey, genus *Cebus* Erxleben 1777 (Cebidae, Primates), from eastern Brazilian Amazonia. *Goeldiana Zoologia* (15):1-3.
- Redford, K. H. and Eisenberg, J. F. 1992. *Mammals of the Neotropics: The Southern Cone, Vol. 2, Chile, Argentina, Uruguay, Paraguay*. Chicago University Press, Chicago.
- Rodríguez-Luna, E. 1993. Avance en la elaboración del Plan de Acción para los Primates de Mesoamérica. *Neotropical Primates* 1(3): 11-13.
- Rosenberger, A. L. 1980. Gradistic views and adaptive radiation of platyrhine primates. *Z. Morph. Anthropol.* 71(2): 157-163.
- Rosenberger, A. L. 1981. Systematics: the higher taxa. In: *Ecology and Behavior of Neotropical Primates*, Vol. 1, A. F. Coimbra-Filho and R. A. Mittermeier (eds.), pp.9-27. Academia Brasileira de Ciências, Rio de Janeiro.
- Rosenberger, A. L. and Coimbra-Filho, A. F. 1984. Morphology, taxonomic status and affinities of the lion tama-

- rins, *Leontopithecus* (Callitrichinae, Cebidae). *Folia Primatol.* 42: 149-179.
- Rosenberger, A. L. and Strier, K. B. 1989. Adaptive radiation of the ateline primates. *J. Hum. Evol.* 18: 717-750.
- Rosenberger, A. L., Setoguchi, T. and Shigehara, N. 1990. The fossil record of callitrichine primates. *J. Hum. Evol.* 19: 209-236.
- Rylands, A. B. 1989. Sympatric Brazilian callitrichids: the black-tufted-ear marmoset, *Callithrix kuhli*, and the golden-headed lion tamarin, *Leontopithecus chrysomelas*. *J. Hum. Evol.* 18(7):679-695.
- Rylands, A. B. 1993. The bare-face tamarins *Saguinus oedipus oedipus* and *Saguinus oedipus geoffroyi*: subspecies or species? *Neotropical Primates*, 1(2): 4-5.
- Rylands, A. B., Spironelo, W. R., Tornisielo, V. L., Lemos de Sa, R. M., Kierulff, M. C. and Santos, I. B. 1988. Primates of the Rio Jequitinhonha valley, Minas Gerais, Brazil. *Primate Conservation* (9): 100-109.
- Rylands, A. B., Santos, I. B. and Mittermeier, R. A. 1991. Distribution and status of *Leontopithecus chrysomelas* in the wild. *Primate Conservation* (12-13). In press.
- Rylands, A. B., Coimbra-Filho, A. F. and Mittermeier, R. A. 1993. Systematics, distributions, and some notes on the conservation status of the Callitrichidae. In: *Marmosets and Tamarins: Systematics, Behaviour and Ecology*, A. B. Rylands (ed.), pp.11-77. Oxford University Press, Oxford.
- Santos, I. B., Mittermeier, R. A., Rylands, A. B. and Valle, C. 1987. The distribution and conservation status of primates in southern Bahia, Brazil. *Primate Conservation* (8):126-142.
- Sanz, V. and Márquez, L. 1994. Conservación del mono capuchino de Margarita (*Cebus apella margaritae*) en la Isla de Margarita, Venezuela. *Neotropical Primates* 2(2): 5-8.
- Seuánez, H. N., Armada, J. L., Freitas, L., R. da Rocha e Silva, Pissinatti, A. and Coimbra-Filho, A. F. 1986. Intraspecific chromosome variation in *Cebus apella* (Cebidae, Platyrhini): the chromosomes of the yellow-breasted capuchin *Cebus apella xanthosternos* Wied, 1820. *Am. J. Primatol.* 10:237-247.
- Schneider, H., Schneider, M. P. C., Sampaio, I., Harada, M. L., Stanhope, M., Czelusniak, J. and Goodman, M. 1993. Molecular phylogeny of the New World monkeys (Platyrhini, Primates). *Molecular Phylogenetics and Evolution* 2(3): 225-242.
- Schneider, M. P. C. Schneider, H., Sampaio, M. I. C., Carvalho-Filho, N. M., Encarnación, F., Montoya, E. and Salzano, F. M. 1995. Biochemical diversity and genetic distances in the Pitheciinae subfamily (Primates, Platyrhini). *Primates* 36(1): 129-134.
- Schneider, H., Sampaio, I., Harada, M. L., Barroso, C. M. L., Schneider, M. P. C., Czelusniak, J. and Goodman, M. In press. Molecular phylogeny of the New World monkeys (Platyrhini, Primates) based on two unlinked nuclear genes: IRBP Intron 1 and -globin sequences. *Am. J. Phys. Anthropol.*
- Silva, B. T. F., Sampaio, M. I. C., Schneider, H., Schneider, M. P. C., Montoya, E., Encarnación, F. and Salzano, F. M. 1992. Natural hybridization between *Saimiri* taxa in the Peruvian Amazonia. *Primates* 33(1): 107-113.
- Silva, B. T. F., Sampaio, M. I. C., Schneider, H., Schneider, M. P. C., Montoya, E., Encarnación, F., Callegari-Jacques, S. M. and Salzano, F. M. 1993. Protein electrophoretic variability in *Saimiri* and the question of its species status. *Am. J. Primatol.* 29: 183-193.
- Silva Jr., J. S. 1988. A range extension for *Saguinus labiatus thomasi*. *Primate Conservation* (9): 23-24.
- Silva Jr., J. S. and Noronha, M. de A. 1995a. A New Record for *Callithrix mauesi* Mittermeier, Schwarz & Ayres, 1992. *Neotropical Primates* 3(3): 79-81.
- Silva Jr., J. S. and Noronha, M. de A. 1995b. On a new species of bare-eared marmoset, genus *Callithrix* Erxleben, 1777, from central Amazonia, Brazil (Primates: Callitrichidae). *Goeldiana Zoologia* (18). In press.
- Silva-López, G., Gill, J. M. and Sanchez-Hernández, A. I. 1995. The Primates of Guatemala: Distribution and Status. Final Report to NYZS The Wildlife Conservation Society, New York. 93pp, January 1995.
- Simons, E. L. 1972. *Primate Evolution: An Introduction to Man's Place in Nature*. Macmillan, New York.
- Simpson, G. G. 1945. The principles of classification and classification of mammals. *Bull. Amer. Mus. Nat. Hist.* 85: 1-350.
- Skinner, C. 1991. Justification for reclassifying Geoffroy's tamarin from *Saguinus oedipus geoffroyi* to *Saguinus geoffroyi*. *Primate Report* 31:77-83.
- Smith, J. D. 1970. The systematic status of the black howler monkey, *Alouatta pigra* Lawrence. *J. Mammal.* 51: 358-369.
- Stallings, J. R. 1985. Distribution and status of primates in Paraguay. *Primate Conservation* (6): 51-58.
- Stanyon, R., Tofanelli, S., Morescalchi, M. A., Agoramoorthy, G., Ryder, O. A. and Wienberg, J. 1995. Cytogenetic analysis shows extensive genomic rearrangements between red howler (*Alouatta seniculus* Linnaeus) subspecies. *Am. J. Primatol.* 35: 171-183.
- Szalay, F. S. and Delson, E. 1979. *Evolutionary History of the Primates*. Academic Press, New York.
- Thorington Jr., R. W. 1985. The taxonomy and distribution of squirrel monkeys (*Saimiri*). In: *Handbook of Squirrel Monkey Research*, L. A. Rosenblum and C. L. Coe (eds.), pp.1-33. Plenum Press, New York.
- Thorington Jr., R. W. 1988. Taxonomic status of *Saguinus tripartitus* (Milne-Edwards, 1878). *Am. J. Primatol.* 15: 367-371.
- Torres de Assumpção, C. 1983. An ecological study of the primates of southeastern Brazil, with a reappraisal of *Cebus apella* races. Ph.D. thesis, University of Edinburgh, Edinburgh.
- Torres, C. 1988. Resultados preliminares de reavaliação das raças do macaco-prego *Cebus apella* (Primates: Cebidae). *Revta. Nordest. Biol.* 6(1):15-28.
- Tyler, D. E. 1991. The evolutionary relationships of *Aotus*. *Folia Primatol.* 56: 50-52.
- Valladares-Padua, C., Padua, S. M. and Cullen Jr., L. 1994a. The conservation biology of the black lion tamarin, *Leontopithecus chrysopygus*: first ten year's report. *Neotropical Primates* 2(suppl.): 36-39.
- Valladares-Padua, C. and Cullen Jr., L. 1994. Distribution, abundance and minimum viable metapopulation of the black lion tamarin *Leontopithecus chrysopygus*. *Dodo*,

- J. Wildl. Preserv. Trusts* 30: 80-88.
- Vieira, C. de C. 1944. Os simios do Estado de São Paulo. *Papéis Avulsos, Zoologia, São Paulo* (4):1-31.
- Vieira, C. de C. 1955. Lista remissiva dos mamíferos do Brasil. *Arq. Zool., São Paulo* 8:341-474.
- Villalba, J. S., Prigioni, C. M. and Sappa, A. C. 1995. Sobre la posible presencia de *Alouatta caraya* en Uruguay. *Neotropical Primates* 3(4). In press.
- Vivo, M. de. 1985. On some monkeys from Rondônia, Brasil (Primates: Callitrichidae, Cebidae). *Pap. Avulsos Zool., São Paulo* 4: 1-31.
- Vivo, M. de. 1991. *Taxonomia de Callithrix Erxleben, 1777* (*Callitrichidae, Primates*). Fundação Biodiversitas, Belo Horizonte.
- Yunis, E. J., Torres de Caballero, O. M. Ramirez, C. and Ramirez, E. 1976. Chromosomal variation in the primate *Alouatta seniculus seniculus*. *Folia Primatol.* 25: 215-224.
- Zunino, G. E., Galliari, C. A. and Colillas, O. J. 1986. Distribución y conservación del mirikina (*Aotus azarae*) en Argentina: resultados preliminares. In: *A Primatología no Brasil - 2*, M. T. de Mello (ed.), pp.305-316. Sociedade Brasileira de Primatologia, Brasília.

## IUCN/SSC PRIMATE SPECIALIST GROUP: NEOTROPICAL SECTION

Chairman: Russell A. Mittermeier

Co-Vice Chairmen for the Neotropical Section: Ernesto Rodríguez Luna and Anthony B. Rylands

### Members

#### ARGENTINA

- *Alejandro Diego Brown* - Laboratorio de Investigaciones Ecológicas de Las Yungas, Universidad Nacional de Tucumán, Casilla de Correo 34, (4107) Yerba Buena, Tucumán.
- *Julio Cesar Ruiz* - Centro Argentino de Primates (CAPRIM), Casilla de Correos 145 (3400) Corrientes
- *Gabriel E. Zunino* - Museo Argentino de Ciencias Naturales "Bernardino Rivadavia", Av. Angel Gallardo 470, Cassilla de Correo 220, Sucursal 5, (1405) Buenos Aires.
- *Marta Mudry* - GIBE, Facultad de Cs. Exactas y Naturales Ciudad Universitaria, Pabellón II, 4º Piso, Av. Cantilo S/N, (1428) Buenos Aires.

#### BELIZE

- *Sharon Matola* - Belize Zoo, P.O. Box 474, Belize City.
- *Robert Horwich* - RD 1, Box 96, Gays Mills, Wisconsin 54631, USA.

#### BOLIVIA

- *Teresa Tarifa Suarez* - Colección Boliviana de Fauna, Casilla 8706, La Paz.
- *Juan Enrique Garcia Yuste* - Asociacion Hombre y Naturaleza, Casilla 218, Santa Cruz de la Sierra.

#### BRASIL

- *Cristina Alves* - Horto-Zoobotânico, Km 21 Rodovia Ilhéus-Itabuna, Caixa Postal 7, Itabuna 45600, Bahia.
- *José Márcio Ayres* - NYZS The Wildlife Conservation Society, Bronx, New York 10460-1099, USA, and Projeto Mamirauá, Departamento de Antropologia, Universidade Federal do Pará, Campus do Guamá, Caixa Postal 531, Belém 66073-250, Pará.
- *Aline P. da Rin Azevedo* - PRIMATAM, Departamento de Zoologia, Museu Paraense Emílio Goeldi, Caixa Postal 399, Belém 66040, Pará.
- *Maria Iolita Bampi* - Departamento de Vida Silvestre, Diretoria de Ecossistemas, Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis (Ibama), SAIN Avenida L4 Norte, Edifício Sede, 70800 Brasília, D.F.
- *Ibsen de Gusmão Câmara* - Avenida das Américas 2300 - C40, Barra da Tijuca, 22640 Rio de Janeiro, Rio de Janeiro.
- *Adelmar F. Coimbra-Filho* - Rua Artur Araripe 60/901, Gávea, 22451-020 Rio de Janeiro, Rio de Janeiro.