

- pp. 440-451. The University of Chicago Press, Chicago.
- Vallejo, A. J. 1995. Estado poblacional, utilización de tipos vegetacionales y crecimiento de *Melanosuchus niger* y *Caiman crocodilus* (Crocodylidae: Alligatorinae) en Zancudococha y Cuyabeno, Amazonía Ecuatoriana. Tesis de Licenciatura, Pontificia Universidad Católica del Ecuador, Quito.
- Whitehead, J. M. 1987. Vocally mediated reciprocity between neighbouring groups of mantled howling monkeys *Alouatta palliata palliata*. *Anim. Behav.* 35: 1615-1627
- Whitehead, J. M. 1995. Vox Alouattinae: A preliminary survey of the acoustic characteristics of long-distance calls of howling monkeys. *Int. J. Primatol.* 16: 121-144.
- Yu-Douglas, W., Hendrickson, T. and Castillo, A. 1997. Ecotourism and conservation in Amazonian Peru: Short-term and long-term challenges. *Environ. Conserv.* 24: 130-138.

DISAPPEARANCE OF INFANTS FOLLOWING MALE TAKEOVERS IN THE BELIZEAN BLACK HOWLER MONKEY (*ALOUATTA PIGRA*)

Robin C. Brockett
Robert H. Horwich
Clara B. Jones

Organisms are expected to employ self-interested tactics and strategies to maximize lifetime probabilities of survival and reproductive success (Trivers, 1985). Behavior programs may differ significantly between the sexes, since selection is thought to operate on the relative parental investment in offspring by males, on the one hand, and females, on the other (Trivers, 1972). Some researchers (e.g., Sugiyama, 1967; Hausfater and Hrdy, 1984) have argued that males may gain a reproductive advantage by killing infants likely to have been sired by non-kin ("infanticide"). This "sexual selection hypothesis" suggests that infanticide shortens a female's interbirth interval through the cessation of lactation and subsequent return of ovarian cycling. Infanticidal males are thought to gain a reproductive advantage by impregnating the dead infant's mother.

Dixon (1998, Table 4.4, p.68) summarizes 48 cases of infanticide observed directly in 13 primate species. Paleotropical species account for 42 of the 48 (88%) cases, and the Hanuman langur (*Presbytis entellus*) accounts for 21 of them (50%). The bias in this database favoring Old World primates, and *P. entellus* in particular, may reflect sampling

error resulting from differential time-investment by researchers. Supporting this idea is the observation that infanticide has been reported most commonly in terrestrial or semi-terrestrial species for which visibility is less of a deterrent to observation.

Infanticide has been reported for four species of *Alouatta* (*A. seniculus*, the red howler monkey: Rudran [1979], Sekulic [1983]; *A. caraya*, the black and brown howler monkey: Zunino *et al.* [1986], Rumiz [1940]; *A. fusca*, the brown howler monkey: Galetti *et al.* [1994]; and *A. palliata*, the mantled howler monkey: Clarke [1981, 1983]). The first three species exhibit polygynous mating systems (after Dixson, 1998) while *A. palliata* groups vary from polygynous to multimale-multifemale (see Crockett and Eisenberg, 1987). Clarke's (1981, 1983) study groups exhibited multimale-multifemale social organization, and infanticide was associated with turnovers in the male hierarchy. Infanticide typically occurs in polygynous (harem or age-graded) or multimale-multifemale mating systems (Dixon, 1988).

We conducted *ad libitum* observations of marked *A. pigra* at the Community Baboon Sanctuary (CBS), Belize. The CBS is a managed reserve formed in 1985 by cooperative agreement among private landowners (Horwich, 1990). Located at 17°33'N, 88°35'W, the CBS is a mosaic of small farms, pastures, and tropical moist forest fragments including riparian habitat along the Belize River (see Horwich and Lyon, 1990). The study area is composed of mapped trails, and >1000 trees have been mapped and identified. Black howlers are generally polygynous with a modal group size of one adult male to several adult females and immatures (Ostro *et al.*, 1999), although multimale groups may be found. Studies of demography, ecology, social organization and behavior are in their early stages (e.g., Horwich, 1983; Silver *et al.*, 1998; Ostro *et al.*, 1999).

As part of a broader study, five incidents of infant disappearance associated with male takeovers were observed (Table 1). These data suggest several topics for further research. First, similar to findings for langurs (*Presbytis* spp.) (Sommer, 1994), there appears to be a male bias in the sex of infants which disappeared. It would be interesting to obtain larger sample sizes in order to evaluate the sex ratios of infants killed, since an offspring's "value" will differ according to its sex and, possibly, the condition of the mother (see Hrdy, 1987). It is possible that infanticide generally occurs in response to some threshold of benefits to costs to the potential victimizer and that the "value" of the po-

Table 1. Observations of infant disappearances at the Community Baboon Sanctuary, Belize. All observations recorded by RCB.

Dates of Takeover	Troop	Male(s) displaced	Displacing male	Mother of infant	Post-takeover copulation observed
2 Feb. - 20 Feb. 1995	Roxie	BBLT, UM	BWB	BBLT BBRT W	yes ¹ no ¹
27 Feb. - 30 Mar. 1995	Baizar	O	BBLT	ORT	no ²
15 Feb. - Mar. 1997	Robin	WLT	Baizar	LLT	yes

¹BBRT gave birth to a male offspring on 13 October 1995 and BBLT to a male offspring on 1 December 1995.

²Copulation attempt observed.

tential victim affects this tradeoff. Some infants may be of such low "value" that they are not worth killing. Alternatively, in some conditions, the costs of infanticide may be prohibitively high. These ideas are supported by the observation that infant disappearance was not observed in association with all male takeovers.

Second, female W died of an intestinal obstruction shortly after the takeover. Although we cannot verify that she was injured during aggressive encounters between displacing and displaced males, it is important to stress that male-male aggression has potential costs (sometimes unexpected) for males that may outweigh the benefits in some conditions.

Third, male BBLT initiated a successful takeover of Baizar troop after he was displaced from Roxie troop by the male BWB. This observation reinforces the benefits of group membership and suggests an apparent cascade effect since we assume that BBLT would not have left his group without the precipitating events of BWB's takeover.

Fourth, our observations show that a single male can take over a multimale group (Roxie troop). Thus, arguments other than superior fighting ability alone must be sought to explain successful takeover strategies. Takeovers should be studied as components of male reproductive strategies, including the role of females in determining which males lead their groups.

Finally, all of our observations occurred in February or March. While this result may be a function of sampling error, systematic studies have been made on a monthly basis at the CBS since 1992, including several thousand contact hours. February and March are the peak of the dry season when preferred food (new leaves, flowers, and fruit) abundance is low and many deciduous trees lose their mature leaves (see Silver, 1998, especially Figs. 2.2 and 2.3), possibly creating stressful conditions for these folivorous monkeys. Food stress in combination with reduced habitat occasioned by deforestation and resultant fragmentation may increase population density and interaction rates, possibly resulting in increased aggressive competition among males (see Kowalewski *et al.*, 1995) and subsequent takeovers. Further research will test the significance of these results.

Our studies suggest that infanticide may be a significant component of a male's lifetime reproductive strategy. Dixon (1998), however, concludes his discussion of infanticide with the statement, "Infanticide is neither *widespread* nor of *general* importance in the evolution of primate social or sexual behaviour." (p.70, italics in the original). He bases this conclusion in part upon the observation that infanticide has been documented in only 12% of primate species. However, Johnstone and Cant (1999), using an evolutionarily stable strategy model, show that the potential for infanticide alone may influence the partitioning of reproduction within groups, even in situations where individuals cannot discriminate their own from another's offspring. Dixon's (1998) conclusion appears premature, awaiting, as is the fate of all speculations, confirmation or disproof

based on future evidence.

Acknowledgements: We are grateful to Dr. K. E. Glander for his assistance in marking the animals. The study was carried out with the support of the National Geographic Society (Grants #5352-94 and #5653-96).

Robin C. Brockett, Zoo Atlanta, Atlanta, Georgia, USA, and Community Conservation Consultants, R.D. 1, Box 96, Gays Mills, Wisconsin 54631, USA, **Robert H. Horwich** Community Conservation Consultants, R.D. 1, Box 96, Gays Mills, Wisconsin 54631, USA, and **Clara B. Jones**, Community Conservation Consultants, and Livingstone College, Department of Psychology, 701 W. Monroe Street, Salisbury, North Carolina 28144, USA.

References

- Clarke, M. R. 1981. Aspects of male behavior in mantled howlers (*Alouatta palliata* Gray) in Costa Rica. *Am. J. Primatol.* 54: 209.
- Clarke, M. R. 1983. Infant killing and infant disappearance following male takeovers in a group of free-ranging howling monkeys (*Alouatta palliata*) in Costa Rica. *Am. J. Primatol.* 56: 241-247.
- Crockett, C. M. and Eisenberg, J. F. 1987. Howlers: variations in group size and demography. In: *Primate Societies*, B. B. Smuts, D. L. Cheney, R. M. Seyfarth, R. W. Wrangham and T. T. Struhsaker (eds.), pp.54-68. The University of Chicago Press, Chicago.
- Dixon, A. F. 1998. *Primate Sexuality*. Oxford University Press, Oxford.
- Galetti, M., Pedroni, F., and Paschoal, M. 1994. Infanticide in the brown howler monkey, *Alouatta fusca*. *Neotropical Primates* 2:6-7.
- Hausfater, G. and Hrdy, S. B. (eds.) 1984. *Infanticide: Comparative and Evolutionary Perspectives*. Aldine, New York.
- Horwich, R. H. 1983. Breeding behaviors in the black howler monkey (*Alouatta pigra*) of Belize. *Primates* 24:222-230.
- Horwich, R. H. 1990. How to develop a community sanctuary—an experimental approach to the conservation of private lands. *Oryx* 24: 95-1025.
- Horwich, R. H. and Lyon, J. 1990. *A Belizean Rainforest*. Orang-Utan Press, Gays Mills, WI.
- Hrdy, S. B. 1974. Male-male competition and infanticide among the langurs (*Presbytis entellus*) of Abu, Rajasthan. *Folia Primatol.* 22: 19-58.
- Hrdy, S. B. 1987. Sex-biased parental investment among primates and other mammals: A critical evaluation of the Trivers-Willard hypothesis. In: *Child Abuse and Neglect: Biosocial Dimensions*, R. J. Gelles and J. B. Lancaster (eds.), pp.97-147. Aldine de Gruyter, New York.
- Johnstone, R. A. and Cant, M. A. 1999. Reproductive skew and indiscriminate infanticide. *Anim. Behav.* 57:243-249.
- Kowalewski, M., Bravo, S. P. and Zunino, G. E. 1995. Aggression between *Alouatta caraya* males in forest patches in northern Argentina. *Neotropical Primates* 3:179-181.
- Ostro, L. E. T., Silver, S. C., Koontz, F. W., Young, T. P. and Horwich, R. H. 1999. Ranging behavior of translocated and established groups of black howler monkeys *Alouatta pigra* in Belize, Central America. *Biol. Conserv.* 87:181-190.
- Rudran, R. 1979. The demography and social mobility of a red howler (*Alouatta seniculus*) population in Venezuela. In: *Verte-*

- brate Ecology in the Northern Neotropics, J. F. Eisenberg (ed.), pp. 107-126. Smithsonian Institution Press, Washington, D.C.
- Rumiz, D. I. 1990. *Alouatta caraya*: Population density and demography in northern Argentina. *Am. J. Primatol.* 2: 279-294.
- Sekulic, R. 1983. Male relationships and infant deaths in red howler monkeys (*Alouatta seniculus*). *Z. Tierpsychol.* 61: 185-202.
- Silver, S. C. 1998. The feeding ecology of translocated howler monkeys, *Alouatta pigra*, in Belize. Ph.D. dissertation, Fordham University, New York.
- Silver, S. C., Ostro, L. E. T., Yeager, C. P., Koontz, F. W. and Horwich, R. 1998. The feeding ecology of the black howler monkey (*Alouatta pigra*) in Northern Belize. *Am. J. Primatol.* 45: 263-279.
- Sommer, V. 1994. Infanticide among the langurs of Jodhpur: Testing the sexual selection hypothesis with a long-term record. In: *Infanticide and Parental Care*, S. Parmigiani and F. S. von Saal (eds.), pp. 137-154. Harwood Academic Publishers, Langehorne, PA.
- Sugiyama, Y. 1967. Social organization of Hanuman langurs. In: *Social Communication Among Primates*. S. A. Altmann (ed.), pp. 221-236. The University of Chicago Press, Chicago.
- Trivers, R. L. 1972. Parental investment and sexual selection. In: *Sexual Selection and the Descent of Man 1871-1971*, B. Campbell (ed.), pp. 136-179. Aldine, Chicago.
- Trivers, R. L. 1985. *Social Evolution*. Benjamin/Cummings Publishing Co., Inc., Menlo Park, CA.
- Zunino, G. E., Chalukian, S. C., and Rumiz, D. I. 1986. Infanticídio e desaparecimento de infantes associados ao reemplazo de machos em grupos de *Alouatta caraya*. In: *A Primatologia no Brasil - 2*, M. T. de Mello (ed.), pp. 185-190. Sociedade Brasileira de Primatologia, Brasília.

News

A NEW SPECIES OF TITI MONKEY IN NORTHEAST BRAZIL

A new species of titi monkey, *Callicebus*, has been described from the Atlantic forest along the coast of the state of Sergipe, northeastern Brazil, by Shuji Kobayashi (Chukyo Women's University, Japan) and Alfredo Langguth (Federal University of Paraíba, Brazil). The species has been named *Callicebus coimbrai*, in honor of Ademar F. Coimbra-Filho, founder and ex-Director of the Rio de Janeiro Primate Center (CPRJ/FEEMA), in recognition of his research and valuable contributions to the biology and conservation of Brazilian primates.

The holotype (an adult female, UFPB1599, in the mammal collection of the Departamento de Sistemática e Ecologia, Universidade Federal da Paraíba) was collected by the authors on 30 January 1994. Two specimens had been shot by a local hunter. The type locality is given as "Proximity of the small village Aragão, in the region of Santana dos Frades about 11.0 km SW of Pacatuba, state of Sergipe, Brazil (GPS 10°32' S, 36°41' W, alt. 90 m)". The locality is south of the Rio São Francisco. The species has been recorded from

between the Rio São Francisco and the Rio Real, which marks the southern border of Sergipe. Although believed to be restricted to the south of the Rio São Francisco, the southern limits to its range have not been established. Titi monkeys have been reported from the south of the Rio Real. Oliver and Santos (1991), for example, obtained reports of titi monkeys from the vicinities of Cachoeira do Abadia and Jandaira, in the north-east of the state of Bahia, which may have been *C. coimbrai*. The western limits to its range are also unknown, but *C. coimbrai* is believed to be restricted to the humid coastal Atlantic forest, and titis inland, in drier, more seasonal formations probably belong to the form *barbarabrownae* Hershkovitz 1990 (see Marinho-Filho and Veríssimo, 1997). Further surveys are urgently needed to establish the limits of the range of this titi, which was listed as "Critically Endangered" by Rylands *et al.* (1995, 1997) even before its description, due to its minute distribution and the widespread destruction of the forests of the region.

C. coimbrai is distinguished from other Atlantic forest titi monkeys by its black forehead, crown and ears, and the buffy trunk. It has pale fur along its sidewhiskers and cheeks, and along the back of the head and nape. The hands and feet are blackish, its tail is orange, and there is a zebra-like striped pattern on the anterior half of the back. Kobayashi and Langguth (1999) also describe distinct features of its dental morphology and craniometry.

On the basis of their study of this new species, along with previous craniometric studies by Kobayashi (1995), Kobayashi and Langguth (1999) argue that the titis of the Atlantic forest should be considered species rather than subspecies of *C. personatus*. They list *C. nigrifrons*, *C. personatus*, *C. melanochir*, and *C. barbarabrownae* besides *C. coimbrai*. Accepting this classification, the Atlantic forest has 21 species of primates, 16 (76%) of which are endemic, and 16 considered threatened according to the 1996 IUCN Red List of Threatened Animals (IUCN, 1996).

Shuji Kobayashi, Department of Asian Studies, Chukyo Women's University, Nadakayama 55, Yokone-cho, Aichi 474-0011, Japan, and **Alfredo Langguth**, Departamento de Sistemática e Ecologia, Universidade Federal da Paraíba, 58059-900 João Pessoa, Paraíba, Brazil.

References

- Hershkovitz, P. 1990. Titis, New World monkeys of the genus *Callicebus* (Cebidae, Platyrrhini): a preliminary taxonomic review. *Fieldiana, Zoology, New Series* (55): 1-109.
- IUCN. 1996. *1996 IUCN Red List of Threatened Animals*. The World Conservation Union (IUCN), Gland, World Conservation Monitoring Center (WCMC), Cambridge.
- Kobayashi, S. 1995. A phylogenetic study of titi monkeys, genus *Callicebus*, based on cranial measurements: I. Phyletic groups of *Callicebus*. *Primates* 36(1): 101-120.
- Kobayashi, S. and Langguth, A. 1999. A new species of titi monkey, *Callicebus* Thomas, from north-eastern Brazil (Primates, Cebidae). *Rev. Bras. Zool.* 16(2): 531-551.
- Marinho-Filho, J. and Veríssimo, E. W. 1997. The rediscovery